

Manuscript Number: IJPara17_387R1

Title: The relationships between faecal egg counts and gut microbial composition in UK Thoroughbreds infected by cyathostomins

Article Type: Full Length Article

Keywords: Helminth-microbiota interactions; Thoroughbred horses; cyathostomins; 16S rRNA sequencing; Adlercreuzia; Methanomicrobia; TM7.

Corresponding Author: Dr. Cinzia Cantacessi,

Corresponding Author's Institution:

First Author: Laura Peachey

Order of Authors: Laura Peachey; Rebecca Molena; Timothy Jenkins; Angela Di Cesare; Donato Traversa; Jane Hodgkinson; Cinzia Cantacessi

Manuscript Region of Origin: UNITED KINGDOM

Abstract: A growing body of evidence, particularly in humans and rodents, supports the existence of a complex network of interactions occurring between gastrointestinal (GI) helminth parasites and the gut commensal bacteria, with substantial effects on both host immunity and metabolic potential. However, little is known of the fundamental biology of such interactions in other animal species. Nonetheless, given the considerable economic losses associated with GI parasites, particularly in livestock and equines, as well as the global threat of emerging anthelmintic resistance, further explorations of the complexities of host-helminth-microbiota interactions in these species are needed. This study characterises, for the first time, the composition of the equine bacterial commensal flora associated with the presence, in faecal samples, of low (Clow) - and high (Chigh) numbers of eggs of an important group of equine GI parasites (i.e. the cyathostomins), prior to and following anthelmintic treatment. High-throughput sequencing of microbial 16S rRNA amplicons and associated bioinformatics and statistical analyses of sequence data revealed strong clustering according to faecal egg counts (FEC) ($P = 0.003$). A trend towards increased populations of Methanomicrobia (class) and Dehalobacterium (genus) was observed in Clow in comparison to Chigh. Anthelmintic treatment in Chigh was associated with a significant reduction of the bacterial phylum TM7 14 days post-ivermectin administration, as well as a transient expansion of Adlercreuzia spp. at 2 days post-treatment. This study provides a first insight into the discovery of the intimate mechanisms governing host-parasite-microbiota interactions in equids, and sets a basis for the development of novel, biology-based intervention strategies against equine GI helminths based on the manipulation of the commensal gut flora.



21 November 2017

Dear Professor Slapeta,

REJOINDER:

R.E. Ms. No. IJPara17_387

Entitled 'The relationships between faecal egg counts and gut microbial composition in UK Thoroughbreds infected by cyathostomins'

We are very grateful for the reviewers' and editor's time and efforts in evaluating this manuscript. We thank the reviewers for their positive comments regarding the quality of our manuscript. The points raised by the reviewers are constructive and have contributed to enhance the manuscript. We have provided responses to individual points in the following rejoinder:

Editor's comment:

Both reviewers acknowledge the value and rigor of the submission. I would only add that you may consider including any data such as QIIME input files inc. metadata and OTUs into Mendeley Data <https://data.mendeley.com/> to enable streamlined reproducibility. Mendeley Data will be directly linked to IJP site upon acceptance. If you do so include the DOI in the data availability section of your Materials & Methods

Our comment: The authors wish to thank the Editor for his kind evaluation. As suggested, raw and curated data have been deposited in the Mendeley database under doi:10.17632/g7chkjrp8f.1. Details have been included in the revised version of the manuscript.

Reviewer #1

This manuscript is an important contribution to the so-far neglected field of the equine microbiome and its relationship with the ubiquitous cyathostomins. This area is likely to provide fruitful in terms of advancing our understanding of basic gut biology, in horses and through a one health approach, in improving control of cyathostomins, including of acute larval cyathostominosis, and in providing improved treatments of colic, colitis and equine inflammatory bowel disease.

The authors have adopted an approach which does not require experimental infection, which is ethically laudable. This approach, however, will require confirmation in future studies from this and other groups in order to demonstrate reproducibility and consistency. Notably, the target group were pregnant broodmares, which in itself may have a bearing on the results.

Our comment: The authors thank Reviewer 1 for their kind comments regarding the overall quality and interest of this manuscript. We agree that due to the experimental approach, repetition of this study in different groups of horses is required to validate our results, as has been noted in the discussion (lines 467-470).

1) There are a few clarifications which the authors may wish to make in order to improve the reader experience. I note the samples were voided naturally. Were any precautions taken to avoid contamination from bacteria on the ground, such as taking samples from the inside of the mass only? Also, were the samples stored /transported anaerobically to preserve the detection of strict anaerobes?

Our comment: We can confirm that the samples were indeed taken from several points inside the faecal mass to avoid contamination from bacteria on the ground, and additionally, that the samples were snap frozen before transport and subsequent storage to preserve detection of bacterial taxa present at the time of voiding. Please see lines 136 and 141, respectively, for clarification of these points.

2) Is there any information on the stage/range of gestation stages of the broodmares in the study?

Our comment: All broodmares enrolled in our study were between months 5-8 of gestation at the start of the trial. This information has been added to the revised manuscript (lines 111-112).

3) The purpose of taking the control samples from non-pregnant animals could perhaps be better explained.

Our comment: Samples were taken from non-pregnant animals to verify that there were no marked alterations to the GI microbiota due to pregnancy which may have affected the outcome of this experiment. Although we cannot unequivocally rule out an interaction between pregnancy and the GI microbiota here, our data suggests that the non-pregnant control animals had a similar microbial profile to the pregnant cohort. Considerations on the possible impact of pregnancy on the composition of the gut microbiota have been added to the revised manuscript (line 138).

4) I don't understand the percentages given for detection of each of the cyathostomin species - please clarify.

Our comment: These percentages represent the proportion of study participants which were positive for any given cyathostomin species. This information has been added to the revised manuscript (lines 215-216).

Reviewer #2

The manuscript is meritorious in terms of the quality, experimental design and methods. Interesting paper and it is worth publishing.

Our comment: We thank Reviewer 2 for their kind comments regarding the quality of our manuscript.

1) In the reference list, Nielsen et al 2010 is reported twice (see lines 690 and 697), delete that to line 697

Our comment: Addressed (line 686).

In conclusion, we consider that we have addressed each of the points raised by each of the reviewers. We agree with the points raised, and modified the text accordingly. The criticisms, suggestions, statements and comments have led to an enhanced manuscript, which we consider to meet the standard for publication in *International Journal of Parasitology*.

Yours sincerely,
Cinzia Cantacessi
On behalf of all authors.



21 November 2017

Dear Professor Slapeta,

REJOINDER:

R.E. Ms. No. IJPara17_387

Entitled 'The relationships between faecal egg counts and gut microbial composition in UK Thoroughbreds infected by cyathostomins'

We are very grateful for the reviewers' and editor's time and efforts in evaluating this manuscript. We thank the reviewers for their positive comments regarding the quality of our manuscript. The points raised by the reviewers are constructive and have contributed to enhance the manuscript. We have provided responses to individual points in the following rejoinder:

Editor's comment:

Both reviewers acknowledge the value and rigor of the submission. I would only add that you may consider including any data such as QIIME input files inc. metadata and OTUs into Mendeley Data <https://data.mendeley.com/> to enable streamlined reproducibility. Mendeley Data will be directly linked to IJP site upon acceptance. If you do so include the DOI in the data availability section of your Materials & Methods

Our comment: The authors wish to thank the Editor for his kind evaluation. As suggested, raw and curated data have been deposited in the Mendeley database under doi:10.17632/g7chkjrp8f.1. Details have been included in the revised version of the manuscript.

Reviewer #1

This manuscript is an important contribution to the so-far neglected field of the equine microbiome and its relationship with the ubiquitous cyathostomins. This area is likely to provide fruitful in terms of advancing our understanding of basic gut biology, in horses and through a one health approach, in improving control of cyathostomins, including of acute larval cyathostominosis, and in providing improved treatments of colic, colitis and equine inflammatory bowel disease.

The authors have adopted an approach which does not require experimental infection, which is ethically laudable. This approach, however, will require confirmation in future studies from this and other groups in order to demonstrate reproducibility and consistency. Notably, the target group were pregnant broodmares, which in itself may have a bearing on the results.

Our comment: The authors thank Reviewer 1 for their kind comments regarding the overall quality and interest of this manuscript. We agree that due to the experimental approach, repetition of this study in different groups of horses is required to validate our results, as has been noted in the discussion (lines 467-470).

1) There are a few clarifications which the authors may wish to make in order to improve the reader experience. I note the samples were voided naturally. Were any precautions taken to avoid contamination from bacteria on the ground, such as taking samples from the inside of the mass only? Also, were the samples stored /transported anaerobically to preserve the detection of strict anaerobes?

Our comment: We can confirm that the samples were indeed taken from several points inside the faecal mass to avoid contamination from bacteria on the ground, and additionally, that the samples were snap frozen before transport and subsequent storage to preserve detection of bacterial taxa present at the time of voiding. Please see lines 136 and 141, respectively, for clarification of these points.

2) Is there any information on the stage/range of gestation stages of the broodmares in the study?

Our comment: All broodmares enrolled in our study were between months 5-8 of gestation at the start of the trial. This information has been added to the revised manuscript (lines 111-112).

3) The purpose of taking the control samples from non-pregnant animals could perhaps be better explained.

Our comment: Samples were taken from non-pregnant animals to verify that there were no marked alterations to the GI microbiota due to pregnancy which may have affected the outcome of this experiment. Although we cannot unequivocally rule out an interaction between pregnancy and the GI microbiota here, our data suggests that the non-pregnant control animals had a similar microbial profile to the pregnant cohort. Considerations on the possible impact of pregnancy on the composition of the gut microbiota have been added to the revised manuscript (line 138).

4) I don't understand the percentages given for detection of each of the cyathostomin species - please clarify.

Our comment: These percentages represent the proportion of study participants which were positive for any given cyathostomin species. This information has been added to the revised manuscript (lines 215-216).

Reviewer #2

The manuscript is meritorious in terms of the quality, experimental design and methods. Interesting paper and it is worth publishing.


Our comment: We thank Reviewer 2 for their kind comments regarding the quality of our manuscript.

1) In the reference list, Nielsen et al 2010 is reported twice (see lines 690 and 697), delete that to line 697

Our comment: Addressed (line 686).

In conclusion, we consider that we have addressed each of the points raised by each of the reviewers. We agree with the points raised, and modified the text accordingly. The criticisms, suggestions, statements and comments have led to an enhanced manuscript, which we consider to meet the standard for publication in *International Journal of Parasitology*.

Yours sincerely,
Cinzia Cantacessi
On behalf of all authors.



Highlights

- We studied qualitative and quantitative differences in faecal microbial profilings between groups of Thoroughbred horses with high- (*Chigh*) and low (*Clow*) faecal egg counts of cyathostomin parasites, pre- and post-anthelmintic treatment.
- A trend towards increased populations of Methanomicrobia (class) and *Dehalobacterium* (genus) was observed in *Clow* in comparison to *Chigh*.
- Anthelmintic treatment in *Chigh* was associated with a significant reduction of the bacterial phylum TM7 14 days post-ivermectin administration, as well as a transient expansion of *Adlercreuzia* spp. at 2 days post-treatment.
- This study sets a basis for the development of novel, biology-based intervention strategies against equine GI helminths based on the manipulation of the commensal gut flora.

**The relationships between faecal egg counts and gut microbial composition in UK
Thoroughbreds infected by cyathostomins**

Peachey L.E.¹, Molena R.A.¹, Jenkins T.P.¹, Di Cesare, A.², Traversa, D.², Hodgkinson J.E.³,
Cantacessi C¹.

¹ Department of Veterinary Medicine, University of Cambridge, Madingley Road, CB3 0ES,
United Kingdom

² Faculty of Veterinary Medicine, University of Teramo, Teramo, 64100, Italy

³ Department of Infection Biology, University of Liverpool, Leahurst, Neston, CH64 7TE,
United Kingdom

Corresponding author: Peachey L.E. lep41@cam.ac.uk; Cantacessi C. cc779@cam.ac.uk

Abstract

A growing body of evidence, particularly in humans and rodents, supports the existence of a complex network of interactions occurring between gastrointestinal (GI) helminth parasites and the gut commensal bacteria, with substantial effects on both host immunity and metabolic potential. However, little is known of the fundamental biology of such interactions in other animal species. Nonetheless, given the considerable economic losses associated with GI parasites, particularly in livestock and equines, as well as the global threat of emerging anthelmintic resistance, further explorations of the complexities of host-helminth-microbiota interactions in these species are needed. This study characterises, for the first time, the composition of the equine bacterial commensal flora associated with the presence, in faecal samples, of low (*Clow*) - and high (*Chigh*) numbers of eggs of an important group of equine GI parasites (i.e. the cyathostomins), prior to and following anthelmintic treatment. High-throughput sequencing of microbial 16S rRNA amplicons and associated bioinformatics and statistical analyses of sequence data revealed strong clustering according to faecal egg counts (FEC) ($P = 0.003$). A trend towards increased populations of Methanomicrobia (class) and *Dehalobacterium* (genus) was observed in *Clow* in comparison to *Chigh*. Anthelmintic treatment in *Chigh* was associated with a significant reduction of the bacterial phylum TM7 14 days post-ivermectin administration, as well as a transient expansion of *Adlercreuzia* spp. at 2 days post-treatment. This study provides a first insight into the discovery of the intimate mechanisms governing host-parasite-microbiota interactions in equids, and sets a basis for the development of novel, biology-based intervention strategies against equine GI helminths based on the manipulation of the commensal gut flora.

Key words: Helminth-microbiota interactions, Thoroughbred horses, cyathostomins, 16S rRNA sequencing, *Adlercreuzia*, Methanomicrobia, TM7.

1. Introduction

Cyathostomins are amongst the most important intestinal nematodes of horses globally (Love et al., 1999; Matthews, 2011; Stratford et al., 2011) with reported prevalence rates as high as 89-100% in equine herds (Mfitilodze and Hutchinson, 1990; Collobert-Laugier et al., 2002; Hinney et al., 2011; Morariu et al., 2016). Clinical signs of cyathostomin infection range from non-specific weight loss to colic and colitis caused by mass emergence of larvae from the large intestinal wall (= larval cyathostominosis), which may prove fatal (Uhlinger, 1991; Murphy and Love, 1997; Lyons et al., 2000; Peregrine et al., 2006). Thoroughbred (TB) young-stock kept in herds are at high risk of developing serious complications of infection, and hence the implementation of effective strategies for parasite control is a top priority for the TB equine industry. Control of cyathostomin infections has traditionally relied on the regular administration of chemotherapeutics (i.e. anthelmintics); however, the frequent and uncontrolled use of these compounds has led to the global emergence of resistant populations of parasites (Nielsen et al., 2014; Peregrine et al., 2014). In particular, foci of multi-drug resistance have been recently reported in UK TB stud farms (Relf et al., 2014). This observation, coupled with the lack of novel anthelmintic compounds licenced for use in equids, represents a ‘Damocle’s sword’ for the UK (and global) equine industry. Therefore, alternative strategies for parasite control are urgently needed; in order to support the discovery of such strategies, a deeper understanding of the complex interactions occurring at the host-parasite interface, particularly at the site/s of infection (i.e. the gut), is required.

While a multitude of factors are responsible for the host-parasite interactions which determine infection outcome, increasing attention is being paid to the complex interplay between GI parasites and the host commensal gut microbiota (Bancroft et al., 2012; Glendinning et al., 2014). Indeed, recent studies have reported significant fluctuations in the composition of the vertebrate gut microbiota associated with helminth infections, that were

accompanied with shifts in both systemic and local immunity (Bancroft et al., 2012; Leung and Loke, 2013; Fricke et al., 2015; Houlden et al., 2015; Cattadori et al., 2016; Gause and Maizels, 2016). However, thus far, knowledge of helminth–microbiota cross-talk relies heavily on studies conducted in humans and/or rodent models of infection and disease (Walk et al., 2010; Rausch et al., 2013; Cantacessi et al., 2014; Lee et al., 2014; Reynolds et al., 2014a; Fricke et al., 2015; Giacomini et al., 2015; Holm et al., 2015; Houlden et al., 2015; Kreisinger et al., 2015; McKenney et al., 2015; Cattadori et al., 2016; Giacomini et al., 2016). Indeed, while only a handful of studies have characterised the composition of the gut microbiota of veterinary species infected by GI helminths thus far (Li et al., 2011; Li et al., 2012; Slapeta et al., 2015; Wu et al., 2012; Duarte et al., 2016; Li et al., 2016; reviewed by Peachey et al., 2017), no data is currently available on the effects of infections by GI helminths, such as cyathostomins, on the composition of the equine commensal flora. Acquiring this fundamental knowledge will be key to the development of novel holistic approaches to equid parasite control, aimed at improving host responses to infections. In this study, we characterise, for the first time, the gut microbial profiles of a cohort of UK TB broodmares with low- and high-numbers of cyathostomin eggs in faeces (as determined by faecal egg counts analysis [FEC]), and examine the effects that administration of a commonly used anthelmintic, i.e. ivermectin, exerts on the overall composition of the gut microbiota as well as relative abundances of individual microbial species.

2. Materials and Methods

2.1. Ethics statement

This study was approved and carried out in strict accordance and compliance with the guidelines of the Institutional Ethical Review Committee, Department of Veterinary Medicine, University of Cambridge, UK (Research Project No. CR190). Written informed consent was obtained from the stud farm from which study samples were collected.

2.2. Sampling and diagnostic procedures

For this study, a cohort of TB broodmares was recruited from a stud farm in the East of England, UK. The stud hosts a group of ~130 pregnant broodmares each year, which are kept at pasture in groups of 2-8 across a 1200-acre farm. All broodmares are subjected to targeted anthelmintic treatments (ivermectin and moxidectin), based on FEC measurements at three monthly intervals. In addition, praziquantel is administered to each broodmare three times a year for tapeworm control, whilst a single moxidectin treatment is administered in late November for control of encysted cyathostomin larvae. Samples used in this study were collected in September-October 2016; all broodmares had received ivermectin and praziquantel in May and August 2016, respectively. A total of 117 TB pregnant broodmares, between 5 and 8 months of gestation at the time of sampling, were screened for infections by cyathostomins. Briefly, individual faecal samples were collected on three consecutive days over a seven-day period; aliquots of each sample were subjected to FEC analysis using a centrifugal floatation technique sensitive to one egg per gram (e.p.g.) (Christie and Jackson, 1982), and to screening for infections by the common equine cestode *Anoplocephala perfoliata* using a double sugar flotation technique (Rehbein et al., 2011). Upon observation of strongyle eggs during FEC analysis, the remaining faecal aliquots were subjected to larval culture to allow for subsequent identification of infecting nematode species using an established Reverse Line Blot (RLB) hybridisation method (Traversa et al., 2007; Cwiklinski et al., 2012). Briefly, genomic DNA was extracted from individual third stage larvae (L3s)

harvested from each larval culture, and the intergenic spacer (IGS) region was amplified by nested PCR using conserved biotin labelled primers (Traversa et al., 2007). The PCR products were then incubated with biodyne C membrane bound specific DNA probes for 21 different cyathostomin species (Cwiklinski et al., 2012), incubated with extravidin peroxidase and visualised using x-ray film. Horses were recruited in our study if they satisfied the following criteria: (i) FEC of >100 e.p.g. (= *Chigh*) or <10 e.p.g. (*Clow*) in three consecutive samples collected over a seven-day period; (ii) matched by approximate age and paddock; (iii) negative for co-infections with other gastrointestinal (GI) helminths; (iv) no antibiotic treatment for at least two months prior to sampling; and (v) no previous anthelmintic treatment other than praziquantel for at least four months prior to sampling. Horses enrolled were kept at pasture for the duration of the study and fed 1kg of custom concentrate mix daily.

2.3. Anthelmintic treatment

Individual, naturally voided, faecal samples were collected from the centre of the faecal mass from *Chigh* and *Clow* animals, as well as from three non-pregnant broodmares on day 0 (D0). Then, an anthelmintic treatment (Eqvalan: ivermectin 0.2 mg/kg) was immediately administered to each animal. Sampling was repeated as above at day 2 (D2) and day 14 (D14) post-treatment (p.t.). A 100g aliquot of each faecal sample was snap frozen, transported to the laboratory and stored at -20 °C within 2h of collection, prior to genomic DNA extraction and sequencing of the bacterial 16S rRNA gene (see below), while the remainder was kept fresh and subjected to FEC analysis as described above.

2.4. High-throughput 16S rRNA sequencing

Genomic DNA was extracted from each faecal sample, as well as from five negative ‘blank’

(= no DNA) controls, using the PowerSoil® DNA Isolation Kit (Qiagen, Carlsbad, CA, USA), according to the manufacturers' instructions. High-throughput sequencing of the V3-V4 hypervariable region of the bacterial 16S rRNA gene was performed on an Illumina MiSeq platform according to the manufacturers' protocols with minor adjustments. Briefly, the V3-V4 region was PCR amplified using universal primers (Forward, 5'-TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG CCT ACG GGN GGC WGC AG-3'; Reverse, 5'-GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA GGA CTA CHV GGG TAT CTA ATC C-3') (Klindworth et al., 2013), that contained the Illumina adapter overhang nucleotide sequences, using the NEBNext® Q5® Hot Start HiFi DNA polymerase (New England Biolabs® Inc, Massachusetts, USA) and the following thermocycling protocol: 98 °C/2 min, followed by 20 cycles of 98 °C/15 s, 63 °C/30 s, and 72 °C/30 s, and a final elongation step of 72 °C/5 min. Amplicons were purified using AMPure XP PCR Purification beads (Beckman Coulter, Brea, California, USA), and the NEBNext hot start high-fidelity DNA polymerase was used for the index PCR with Nextera XT index primers (Illumina, San Diego, California, USA) according to the following thermocycling protocol: 98 °C/30 s, 8 cycles of 98 °C/10 s, 65 °C/75 s and at 65 °C/5 min. The indexed samples were purified using AMPure XP beads, quantified using the Qubit Quant-iT™ dsDNA Broad-Range Assay Kit (Life Technologies, Carlsbad, California, USA), and equal quantities from each sample pooled. The resulting pooled library was quantified using the NEBNext® Library Quant Kit for Illumina® (New England Biolabs® Inc) and sequenced on an Illumina MiSeq platform using the v3 chemistry (301 bp paired-end reads).

2.5. Bioinformatics analyses

Raw paired-end Illumina reads were trimmed for 16S rRNA gene primer sequences using Cutadapt (<https://cutadapt.readthedocs.org/en/stable/>), and reads were joined using the

174 Quantitative Insights Into Microbial Ecology (QIIME) software suite (version 1.9.0)
175 (Caporaso et al., 2010). Successfully joined sequences were quality filtered in QIIME using
176 the ‘usearch_qf’ script with default settings. Then, sequences were clustered into Operational
177 Taxonomic Units (OTUs) on the basis of similarity to known bacterial sequences available in
178 the Greengenes database (v13.8; <http://greengenes.secondgenome.com/>; 97% sequence
179 similarity cut-off) using the UCLUST software; sequences that could not be matched to
180 references in the Greengenes database were clustered *de novo* based on pair-wise sequence
181 identity (97% sequence similarity cut-off). The first selected cluster seed was considered as
182 the representative sequence of each OTU. Then, representative sequences were assigned to
183 taxonomy using the UCLUST software. OTUs assigned to sequences obtained from the no-
184 DNA control samples, as well as singleton OTUs, were removed prior to downstream
185 analysis. Normalisation was carried out by generating a subsampled OTU table by random
186 sampling (without replacement) of the input OTU table using an implementation of the
187 Mersenne twister algorithm (<http://www.numpy.org>). Cumulative-sum scaling (CSS) and
188 log2 transformation were applied to account for the non-normal distribution of taxonomic
189 counts data. Statistical analyses were executed using the Calypso software
190 (cgenome.net/calypso/); samples were clustered using supervised Canonical Correlation
191 Analysis including FEC (*Chigh:Clow*) and time-point (D0, D2 and D14 p.t.) as explanatory
192 variables. Differences in bacterial alpha diversity (Shannon diversity) between groups were
193 evaluated using paired t-test or ANOVA, depending on number of groups for comparison.
194 Beta diversity was calculated using weighted UniFrac distances and based on the matrices,
195 differences in beta diversity were calculated using Permutational Analysis of Multivariate
196 Dispersions (PERMDISP) through the ‘betadisper’ function (Anderson et al., 2006).
197 Differences in the composition of the faecal microbiota between groups were assessed using
198 the LEfSe (Linear discriminant analysis Effect Size) workflow (Segata et al., 2011), by

assigning FEC/timepoint ‘groupings’ as comparison class. All statistical analyses were repeated on a sub-group of horses with FEC >200 e.p.g. (n=8) and 0 e.p.g. (n=7), hereafter referred to as ‘C200’ and ‘C0’, respectively.

3. Results

3.1 Prevalence of cyathostomin infection and group selection

Of 117 broodmares examined for cyathostomin infection, 36 matched the criteria outlined in section 2.2. and were therefore enrolled in this study. Of these, 18 broodmares had an average FEC of ≥ 100 e.p.g. (range 100 - 418), and were thus enrolled as *Chigh* (Table 1). Amongst these, 8 were defined as C200 based on average FEC of ≥ 200 e.p.g. (Table 1). Conversely, 18 broodmares with FEC of ≤ 10 e.p.g. (range 0 - 10) were enrolled as *Clow* and, of these, 7 could be defined as C0 (Table 1). While no known effect size estimates for changes in microbiota due to parasitism in horses are currently available, data from previous studies in other host:helminth systems (effect size: 1.5; cf. Giacomini et al., 2015, 2016) provided us with 84% power to detect changes in gut microbial composition between groups (n=18 in each *Chigh* and *Clow* group) using canonical analysis of principal coordinates.

Larval culture followed by species identification *via* RLB revealed infections by the following cyathostomin species: *Cyathostomum* (*Cya.*) *catinatum* (88%), *Cylicostephanus* (*Cys.*) *longibursatus* (71%), *Cylicocyclus* (*Cyc.*) *nassatus* (59%), *Coronocyclus coronatus* (41%), *Cyc. calicatus* (41%), *Cyc. radiatus* (29%), *Cys. goldi* (24%), *Cys. leptostomus* (18%), *Cya. pateratum* (18%), *Cyc. ashworthi* (18%) and *Cyc. insigne* (12%). FEC analysis performed on samples collected at D2 and D14 p.t. showed FEC reduction rates (FECR) of > 95% in all treated animals (Table 1).

3.2 Microbiota profiling

A total of 39,461,550 raw paired-end reads were generated from DNA faecal extracts of *Chigh* and *Clow* broodmares and subjected to further processing. Following primer trimming, joining of paired-end reads, filtering of low-quality sequences and removal of ‘contaminant’ and singleton OTUs, a total of 8,077,490 (mean = $73,423 \pm 3,610$) high quality sequences were retained for further bioinformatics analysis (not shown). Raw and curated sequence data generated in this study are available from Mendeley Data at doi:10.17632/g7chkjrp8f.1. The rarefaction curves generated following in-silico subtraction of low-quality and contaminant sequences indicated that the vast majority of faecal bacterial communities were represented in the remaining sequence data, thus allowing us to undertake further analyses. These sequences were assigned to 95,286 OTUs and 15 bacterial phyla, respectively. The phyla Bacteroidetes (39.9%) and Firmicutes (34.0%) were predominant in all samples, followed by the phyla Verrucomicrobia (12.0%), Spirochaetes (3.9%), Fibrobacteres (2.4%), Cyanobacteria (1%), Proteobacteria (0.9%), Euryarcheota (0.4%), Tenericutes (0.4%), TM7 (0.3%), Actinobacteria (0.3%), Lentisphaerae (0.3%), Synergistetes (0.2%), WPS-2 (0.2%) and Planctomycetes (0.1%) (Fig. 1), while 3.4% of OTUs could not be assigned to any bacterial group. Predominant sub-taxa were Bacteroidia (class), Bacteroidales (order) and Bacteroidales (family) within the phylum Bacteroidetes, and Clostridia (class), Clostridiales (order) and Ruminococcae (family) within the Firmicutes (Fig. 1). Two samples, LVN1 and HS2, differed markedly in the relative proportions of the two most abundant phyla, Bacteroidetes and Firmicutes, when compared with samples from other broodmares (Fig. 1), likely indicating dysbiosis. Therefore, in order to reduce biases due to these potential ‘outliers’, these samples were excluded from further statistical analyses (Fig. 1).

Figure 1: Bar charts depicting the relative abundances of faecal bacterial phyla from broodmares with faecal egg counts (FEC) >100 eggs per gram (e.p.g.) (= *Chigh*) and <10 e.p.g. (= *Clow*), according to sampling time point (i.e. pre-anthelmintic treatment [D0], and 2 and 14 days post-treatment [D2 and D14, respectively], and in non-pregnant controls (NPC). Samples from broodmares with FEC >200 e.p.g. (*C200*) and 0 (*C0*) are shown in red.

3.3. Differences in microbial composition between Chigh and Clow, and pre- and post-anthelmintic treatment

Microbial community profiles of each sample were grouped by hierarchical clustering and ordinated by supervised CCA. Using these methods, a significant association was observed between microbial composition and FEC (*Chigh versus Clow*) ($P = 0.003$), while clustering according to time point pre- and post- anthelmintic treatment (*D0 versus D14*) did not reach statistical significance ($P = 0.686$) (Fig. 2a). CCA of *C200 versus C0* led to a clear separation according to FEC ($P = 0.001$), whilst the effect of anthelmintic treatment remained insignificant ($P = 0.811$) (Fig. 2b).

Figure 2: The microbial composition of faecal samples ordered by supervised canonical correspondence analysis (CCA) from broodmares with (a) faecal egg counts (FEC) >100 eggs per gram (e.p.g.) (= *Chigh*) and <10 e.p.g. (= *Clow*), pre-anthelmintic treatment [D0] and at 14 days post-treatment [D14], (b) with FEC >200 e.p.g. (*C200*) and 0 (*C0*) at D0 and D14.

No significant differences in OTU alpha diversity (Shannon Index) were recorded between *Chigh* and *Clow*, or between samples collected at D0, D2 and D14 p.t. (Fig.3 a,b,c). A trend

towards increased alpha diversity in *Chigh* versus *Clow* at all time-points was observed ($P = 0.087$) (Fig 3a). This trend was also observed when C200 samples were compared to C0 at D0, despite smaller group sizes ($P = 0.102$) (Fig. 3c). No significant differences in beta diversity, as measured by PERMDISP, were observed between groups (Fig. 4).

Figure 3: Shannon diversity charts comparing faecal microbial alpha diversity of broodmares (a) with faecal egg counts (FEC) >100 eggs per gram (e.p.g.) (= *Chigh*) and <10 e.p.g. (= *Clow*) at all time points (i.e. pre-anthelmintic treatment [D0] and 2 and 14 days post-treatment [D2 and D14, respectively]), (b) *Chigh* and *Clow* at D0 only, (c) with FEC >200 e.p.g. (C200) and 0 (C0) at D0 only, and (d) *Chigh* and (e) *Clow* at D0, D2 and D14.

Figure 4: Permutational Analysis of Multivariate Dispersions (PERMDISP) plots comparing the faecal microbial beta diversity of broodmares (a) with faecal egg counts (FEC) >100 eggs per gram (e.p.g.) (*Chigh*) and <10 e.p.g. (*Clow*) at all time points (i.e. pre-anthelmintic treatment [D0] and 2 and 14 days post-treatment [D2 and D14, respectively]), (b) *Chigh* and *Clow* at D0 only, (c) with FEC >200 e.p.g. (C200) and 0 (C0) at D0 only, and (d) *Chigh* and (e) *Clow* at D0, D2 and D14.

Differences in abundance of individual taxa at the phylum, class, order, family, genus and species level were detected between *Chigh* and *Clow* samples, as well as between samples collected at D0, and D2 and D14 p.t. (Fig. 5). Samples from *Clow* at D0 (pre-treatment) were characterised by an increased abundance of Methanobacteria (class), *Dehalobacterium* (genus) and unclassified *Dehalobacterium* and *Ruminococcus* (species) compared with samples from *Chigh* (Fig. 5a). The same taxa were increased in C0 compared with C200,

with the addition of methanogens of the family Methanocorpusculaceae belonging to order Methanomicrobiales, class Methanobacteria, phylum Euryarchaeota; order Endomicrobiales, phylum Elusimicrobia; Rickettsiales (order, family, genus, species); family Bacteroidaceae, genus BF311 and species RFN20 (Fig. 5b). The taxa GMD14H09 (order, family, genus, species) of the phylum Proteobacteria were increased in samples from C200 compared with C0 (Fig. 5b). Anthelmintic treatment in *Chigh* was accompanied by a decrease in the phylum TM7 at D14, when compared with pre-treatment samples (Fig. 5c). Additionally, the taxa *Adlercreutzia* and R445B were increased at D2 and D14, respectively, compared with D0 samples (Fig. 5c). In *Clow*, treatment was also associated with an increase in R445B (family, genus, species) at D14 (Fig. 5d).

Figure 5: Linear discriminant analysis Effect Size (LEfSe) bar charts depicting differences in abundance of individual bacterial taxa at the phylum, class, order, family, genus and species level in faecal samples from broodmares (a) with faecal egg counts (FEC) >100 eggs per gram (e.p.g.) (*Chigh*) and <10 e.p.g. (*Clow*), (b) with FEC >200 e.p.g. (*C200*) and 0 (*C0*) at D0, and in (c) *Chigh* and (d) *Clow* according to sampling time point (i.e. pre-anthelmintic treatment [D0] and 2 and 14 days post-treatment [D2 and D14, respectively]).

4. Discussion

This study is the first to report a significant association between numbers of cyathostomin eggs in faecal samples from UK Thoroughbreds and the composition of the host gut microbiota. A particularly significant shift in microbial profiles was observed when the faecal bacterial populations of a group of broodmares with FEC of >200 e.p.g. were compared with those with observed FEC of 0. These data are consistent with observations from published

studies in both humans and other veterinary species, including rodent models of infection and disease (Lee et al., 2014; Holm et al., 2015; Houlden et al., 2015; McKenney et al., 2015; Duarte et al., 2016; Li et al., 2016). In addition, the administration of a routinely used anthelmintic (i.e. ivermectin) to both *Chigh* and *Clow* resulted in further progressive changes of the microbial profiling of treated horses. While such changes did not reach statistical significance when analysed using a multivariate model, this trend suggests that parasite-associated modifications in gut microbiota may be transient, and dependent on the presence of live infections, a hypothesis which requires thorough testing.

Overall, the bacterial phyla identified in this study were consistent between groups of animals enrolled in this study; this observation differs from results from previous studies that had reported significant variability in faecal microbial profiling between horses, largely related to variations in diet and age, and presence of underlying diseases (Costa et al., 2012; Daly et al., 2012; Steelman et al., 2012; O' Donnell et al., 2013; Dougal et al., 2014; Fernandes et al., 2014; Weese et al., 2014). Thus, our finding likely indicates that the impact of such confounding factors was successfully minimised by our study design, and that the recorded differences in microbial composition were indeed associated to parasite infections. Bacteria belonging to the phylum Bacteroidetes were predominant in animals examined in our study; conversely, other investigations had reported Firmicutes as being the most prevalent phylum in the horse gut flora (Costa et al., 2012; Shepherd et al., 2012; Dougal et al., 2014; Fernandes et al., 2014; Weese et al., 2014; Costa et al., 2015a; Costa et al., 2015b; Proudman et al., 2015). Dietary differences between horse cohorts enrolled in this and previous studies are likely to be responsible for this discrepancy (cf. Daly et al., 2012; Fernandes et al., 2014).

Overall, a trend towards increased microbial alpha diversity, i.e. the number of different OTUs in each sample [‘richness’] and their relative abundance [= ‘evenness’] (Tuomisto, 2010), was observed in samples from *Chigh* compared with those from *Clow* at D0 (pre-anthelmintic treatment) and in *C200 versus C0*, although these differences did not reach statistical significance. Nevertheless, this observation is supported by the results of a number of previous studies in other host:helminth systems, in which the establishment of parasite infections was associated with an overall increase in alpha diversity of the gut microbiota (Broadhurst et al., 2012; Lee et al., 2014; Giacomini et al., 2015; Giacomini et al., 2016). Given that a number of inflammatory GI and/or systemic diseases are accompanied by a reduced alpha diversity (Manichanh et al., 2006; Sepehri et al., 2007; Abrahamsson et al., 2012, 2014), the increase in GI microbial diversity observed in the presence of helminth infections has been hypothesized to represent one of the possible mechanisms by which parasites suppress host inflammatory responses, thus ensuring their long-term survival in the host gut (Bancroft et al., 2012; Glendinning et al., 2014). Therefore, the trends towards increased alpha diversity observed in the faecal microbiota of horses moderately infected by cyathostomins may also result from an increase in gut homeostasis promoted by the parasite. Future studies evaluating the prevalence and incidence of equine inflammatory diseases (e.g. inflammatory bowel disease and recurrent airway obstruction) in the presence or absence of parasite infections could represent significant first steps in this area of research.

In addition to global microbial diversity, significant variations in the abundance of specific bacterial taxa were observed between groups. In particular, a higher abundance of microorganisms belonging to the class Methanomicrobia was observed in *Clow* (D0) when compared to *Chigh*. This difference was exacerbated in *C200 versus C0*, with further significant increases in methanogens belonging to class Methanomicrobia recorded in *C0*,

thus suggesting a negative correlation between methanogen abundance and FEC. Methanomicrobia belong to the phylum Euryarchaeota, kingdom Archaea and are phylogenetically distinct from bacteria and eukaryotes, although they retain the prokaryote 16S rRNA gene (Woese and Gupta, 1981; Winker and Woese, 1991). Particularly in ruminants, the role of the Archaeal methanogens in the digestion of fibre has been well documented (St-Pierre et al., 2015). In equids, little is known of the functional diversity of methanogens; however, consistent with our findings, a recent study reported Methanomicrobiales as being predominant in the horse gut (Lwin and Matsui, 2014). The underlying mechanisms by which GI helminths may be promoting a reduction in populations of methanogens are yet unclear. Similarly to hypotheses formulated for other host:helminth systems, cyathostomins may prevent expansion of methanogens directly, e.g. *via* their excretory secretory products, or indirectly *via* parasite-induced changes in mucosal immunity (reviewed by Peachey et al., 2017). Alternatively, a high abundance of methanogens prior to helminth infections may bias host immune responses against cyathostomins, thus resulting in the observed low (or absent) parasite burdens. Interestingly, some methanogens (i.e. *Methanosphaera stadtmanae*) have been shown to regulate Th17 responses in mice (Blais Lecours et al., 2011; Bernatchez et al., 2017); in turn, these responses have been linked to the ability of mice to clear experimental infections by *Heligmosomoides polygyrus* (Reynolds et al., 2014b). Mechanistic studies aimed to evaluate the effects of expanding populations of gut methanogens on host mucosal responses and, in turn, GI helminth establishment, may help elucidating these interactions.

An increased abundance of Methanomicrobia in *C*_{low} and *C*₀ may also be linked to other environmental factors that are simultaneously responsible for the low FEC observed. An example is represented by the horse grazing behaviour; indeed, it is known that some

individuals within a herd favour less nutritional swards of grass in order to avoid faecal contamination (Hutchings et al., 2000). In turn, as animal faeces often act as fertilisers, as a result, individuals favouring nutritious grass are exposed to higher numbers of infective larvae. Grazing different swards of grass may also impact on dietary fibre levels, and thus on gut methanogen populations, as observed in ruminants (McAllister et al., 1996). In horses, dietary factors have also been associated with changes in abundance of Methanomicrobia; for example, *Methanocorpusculum* archaea were observed at a median of 17.7% in horses fed a forage-grain diet, and at a median of 31.9% in horses maintained on pasture (Fernandes et al., 2014). Differences in grazing behaviour between individuals may also be accountable for the increased abundance of bacteria of the phylum Elusimicrobia in C0 versus C200 as these taxa are primarily a component of termite hind-gut microbiota (Gómez and González-Megías, 2007; van Klink et al., 2015; Mikaelyan et al., 2017). Experimental cyathostomin infections of stabled horses may eliminate the effect of grazing behaviour on gut microbial profiles, although ethical concerns may prevent the execution of such a study in the future.

In contrast to uninfected horses, the faecal microbial profiles of C200 were characterised by an increased abundance of GMD14H09, phylum Proteobacteria, class Deltaproteobacteria. Increases in Proteobacteria abundance have repeatedly been reported in association with helminth infections, e.g. in mice infected by *Trichuris muris* and *H. polygyrus*, pigs infected by *T. suis*, and rabbits infected by *Trichostrongylus retortaeformis* (Li et al., 2012; Holm et al., 2015; Cattadori et al., 2016). Proteobacteria are known increase in the presence of GI inflammation (Shin et al., 2015); hence, the expansion of populations of Proteobacteria in the faecal microbiota of horses with higher infection burdens may be indicative of an inflammatory status of the intestinal tract of these horses at the time of sampling.

One of the objectives of this study was to assess the impact of anthelmintic treatment on the faecal microbial profiling of cyathostomin-infected horses. In particular, ivermectin administration to *Chigh* was followed by a significant decrease in populations of the phylum TM7 at D14. Since the relative abundance of TM7 remained unchanged following ivermectin administration in *Clow*, it is tempting to speculate that a mutualistic association may exist between TM7 and cyathostomins, whereby each promote establishment of the other, similarly to the mutual relationship described for Lactobacillaceae and *H. polygyrus* (Reynolds et al., 2014a). Bacteria belonging to the phylum TM7 are obligate epibionts of *Actinomyces* spp. (He et al., 2015), and are thus uncultivable. While TM7 have not previously been linked to GI helminth infections, this phylum of bacteria has been associated with mucosal inflammatory disease in humans (Kuehbachner et al., 2008). Interestingly, TM7 isolates have been shown to repress the induction of TNF- α production in macrophages infected by *Actinomyces odontolyticus*, thus suggesting a potential immune suppressive activity (He et al., 2015); hence, TM7 may promote the establishment of cyathostomins by suppressing effective anti-parasite immune responses. Furthermore, an increase of the taxa *Adlercreutzia* (phylum actinobacteria) and R445B (phylum Lentisphearae) was observed in *Chigh* at D2 and D14, respectively. The latter was also increased in *Clow* at D14, suggesting that this change was unrelated to cyathostomin removal. Bacteria of the genus *Adlercreutzia* produce the metabolite equol (Maruo et al., 2008), a known anti-inflammatory and vasodilator (Blay et al., 2010). Thus, it could be hypothesised that increases in populations of *Adlercreutzia* and its metabolites following ivermectin administration might contribute to the emergence of hypobiotic larval stages of cyathostomins (which is known to occur post-anthelmintic treatment; Lyons et al., 2000), via the suppression of effective mucosal immune responses. This hypothesis requires testing in controlled mechanistic experiments.

FEC are often utilised as proxy of parasite infection burdens; however, several investigations

have confuted this practice, as weak correlations have been detected between FEC and parasite burdens in horses with >500 e.p.g. of faeces (Nielsen et al., 2010). While the FEC cut-offs used in this study are indicative of differing infection burdens between groups, any inference on the relationships between number of worms in the horse intestine and gut microbial profiling must be taken with caution. Ethical considerations prevent us from performing post-mortem total worm counts on experimentally infected horses; nevertheless, in the future, it may be possible to establish unequivocal relationships between cyathostomin infection burdens (including encysted larvae) and gut microbial profiling from samples collected in abattoir.

Clearly, a complex network of host-parasite interactions, as well as environmental factors, contribute to the findings reported in this study, and thus further work is needed to disentangle the causality of these relationships. However, one key question that needs addressing is whether differences in host immunity may be associated with significant changes in gut microbial composition (and *vice versa*) and, if such is the case, whether the horse gut microbiota could be manipulated to improve resistance to helminth infection. Indeed, previous investigations in cattle and mice have reported that host genes encoding for antimicrobial proteins are up-regulated in the mucosae of animals resistant to helminth infection (D'Elia et al., 2009; Li et al., 2015). In addition, dietary supplementation with both pro- (Bautista-Garfias et al., 1999; Bautista-Garfias et al., 2001; Martinez-Gomez et al., 2009; Martinez-Gomez et al., 2011; Oliveira-Sequeira et al., 2014; El Temsahy et al., 2015) and pre-biotics (Petkevicius et al., 2003; Petkevicius et al., 2004; Thomsen et al., 2005; Petkevicius et al., 2007; Jensen et al., 2011), has led to significant reductions in worm burdens in murine and swine helminth infection models, thus indicating that alterations to the bacterial flora in the host GI tract may bias host immune responses against parasites. Further

characterisation of equine host mucosal responses and GI microbiota, in the presence or absence of helminth infection and accompanied by total enumeration of infecting parasites, is a key area of future research, as it may lead to the identification of microbial factors linked to host susceptibility.

5. Conclusion

Cyathostomin infection in horses was associated with global shifts in faecal microbial composition and diversity, in accordance with previous studies in other host: helminth systems, as well as significant changes in specific populations of gut bacteria. Such changes predominantly involved ‘minor’ phyla, thus suggesting that the equine ‘core gut microbiota’ remains unaltered in the presence of burdens of cyathostomins such as those observed in this study. Our findings also support the hypothesis that selected bacterial taxa, and/or their metabolites, may play roles in biasing the host immune response either for, e.g. TM7, or against, e.g. Methanomicrobia, cyathostomin infection in horses. These data pave the way for future mechanistic studies aimed to identify microbial factors linked to host susceptibility, and to manipulate the GI microbiota of horses (e.g. via dietary or probiotic interventions), in order to improve resistance to cyathostomins.

Role of the Funding Body

L.E.P. is a Post-Doctoral Research Fellow co-funded by the Horserace Betting Levy Board, the Thoroughbred Breeders' Association and the British European Breeders Fund (Grant no. VET/EPDF/2016-2). T.P.J. is the grateful recipient of a Ph.D. scholarship by the Biotechnology and Biological Sciences Research Council. Research in C.C. laboratory is funded by grants by the Royal Society and the Isaac Newton Trust/ Wellcome Trust/ University of Cambridge.

Acknowledgements

The authors are very grateful for the cooperation and assistance provided by the management and staff of the Thoroughbred stud farm where this study was performed. Additionally, the authors would like to thank Professor James Wood (University of Cambridge) for constructive comments.

References

- Abrahamsson, T.R., Jakobsson, H.E., Andersson, A.F., Bjorksten, B., Engstrand, L., Jenmalm, M.C., 2012. Low diversity of the gut microbiota in infants with atopic eczema. *J. Allergy Clin. Immunol.* 129, 434-440, 440 e431-432.
- Abrahamsson, T.R., Jakobsson, H.E., Andersson, A.F., Bjorksten, B., Engstrand, L., Jenmalm, M.C., 2014. Low gut microbiota diversity in early infancy precedes asthma at school age. *Clin. Exp. Allergy.* 44, 842-850.
- Anderson, M.J., Ellingsen, K.E., McArdle, B.H., 2006. Multivariate dispersion as a measure of beta diversity. *Ecol. Lett.* 9, 683-693.
- Bancroft, A.J., Hayes, K.S., Grenis, R.K., 2012. Life on the edge: the balance between macrofauna, microflora and host immunity. *Trends Parasitol.* 28, 93-98.

515 Bautista-Garfias, C.R., Ixta, O., Orduna, M., Martinez, F., Aguilar, B., Cortes, A., 1999.
 516 Enhancement of resistance in mice treated with *Lactobacillus casei*: effect on *Trichinella*
 517 *spiralis* infection. Vet. Parasitol. 80, 251-260.

518 Bautista-Garfias, C.R., Ixta-Rodriguez, O., Martinez-Gomez, F., Lopez, M.G., Aguilar-
 519 Figueroa, B.R., 2001. Effect of viable or dead *Lactobacillus casei* organisms administered
 520 orally to mice on resistance against *Trichinella spiralis* infection. Parasite. 8, S226-228.

521 Bernatchez, E., Gold, M.J., Langlois, A., Blais-Lecours, P., Boucher, M., Duchaine, C.,
 522 Marsolais, D., McNagny, K.M., Blanchet, M.R., 2017. *Methanosphaera stadtmanae* induces
 523 a type IV hypersensitivity response in a mouse model of airway inflammation. Physiol. Rep.
 524 5.

525 Blais Lecours, P., Duchaine, C., Taillefer, M., Tremblay, C., Veillette, M., Cormier, Y.,
 526 Marsolais, D., 2011. Immunogenic properties of archaeal species found in bioaerosols. PLoS
 527 One. 6, e23326.

528 Blay, M., Espinel, A.E., Delgado, M.A., Baiges, I., Blade, C., Arola, L., Salvado, J., 2010.
 529 Isoflavone effect on gene expression profile and biomarkers of inflammation. J. Pharm.
 530 Biomed. Anal. 51, 382-390.

531 Broadhurst, M.J., Ardesir, A., Kanwar, B., Mirpuri, J., Gundra, U.M., Leung, J.M., Wiens,
 532 K.E., Vujkovic-Cvijin, I., Kim, C.C., Yarovsky, F., Lerche, N.W., McCune, J.M., Loke, P.,
 533 2012. Therapeutic helminth infection of macaques with idiopathic chronic diarrhea alters the
 534 inflammatory signature and mucosal microbiota of the colon. PLoS Pathog. 8, e1003000.

535 Cantacessi, C., Giacomini, P., Croese, J., Zakrzewski, M., Sotillo, J., McCann, L., Nolan,
 536 M.J., Mitreva, M., Krause, L., Loukas, A., 2014. Impact of experimental hookworm infection
 537 on the human gut microbiota. J. Infect. Dis. 210, 1431-1434.

538 Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K.,
 539 Fierer, N., Pena, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T., Knights, D.,

540 Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D., Pirrung, M., Reeder,
 541 J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A., Widmann, J., Yatsunenko, T., Zaneveld,
 542 J., Knight, R., 2010. QIIME allows analysis of high-throughput community sequencing data.
 543 Nat. Methods. 7, 335-336.

544 Cattadori, I.M., Sebastian, A., Hao, H., Katani, R., Albert, I., Eilertson, K.E., Kapur, V.,
 545 Pathak, A., Mitchell, S., 2016. Impact of helminth infections and nutritional constraints on
 546 the small intestine microbiota. PLoS One. 11, e0159770.

547 Christie, M., Jackson, F., 1982. Specific identification of strongyle eggs in small samples of
 548 sheep faeces. Res. Vet. Sci. 32, 113-117.

549 Collobert-Laugier, C., Hoste, H., Sevin, C., Dorchies, P., 2002. Prevalence, abundance and
 550 site distribution of equine small strongyles in Normandy, France. Vet. Parasitol. 110, 77-83.

551 Costa, M.C., Arroyo, L.G., Allen-Vercoe, E., Stampfli, H.R., Kim, P.T., Sturgeon, A.,
 552 Weese, J.S., 2012. Comparison of the fecal microbiota of healthy horses and horses with
 553 colitis by high throughput sequencing of the V3-V5 region of the 16S rRNA gene. PLoS One.
 554 7, e41484.

555 Costa, M.C., Silva, G., Ramos, R.V., Staempfli, H.R., Arroyo, L.G., Kim, P., Weese, J.S.,
 556 2015a. Characterization and comparison of the bacterial microbiota in different
 557 gastrointestinal tract compartments in horses. Vet. J. 205, 74-80.

558 Costa, M.C., Stampfli, H.R., Arroyo, L.G., Allen-Vercoe, E., Gomes, R.G., Weese, J.S.,
 559 2015b. Changes in the equine fecal microbiota associated with the use of systemic
 560 antimicrobial drugs. BMC Vet. Res. 11, 19.

561 Cwiklinski, K., Kooyman, F.N., Van Doorn, D.C., Matthews, J.B., Hodgkinson, J.E., 2012.
 562 New insights into sequence variation in the IGS region of 21 cyathostomin species and the
 563 implication for molecular identification. Parasitology. 139, 1063-1073.

564 D'Elia, R., DeSchoolmeester, M.L., Zeef, L.A., Wright, S.H., Pemberton, A.D., Else, K.J.,
 565 2009. Expulsion of *Trichuris muris* is associated with increased expression of angiogenin 4 in
 566 the gut and increased acidity of mucins within the goblet cell. BMC Genomics. 10, 492.
 567 Daly, K., Proudman, C.J., Duncan, S.H., Flint, H.J., Dyer, J., Shirazi-Beechey, S.P., 2012.
 568 Alterations in microbiota and fermentation products in equine large intestine in response to
 569 dietary variation and intestinal disease. Br. J. Nutr. 107, 989-995.
 570 Dougal, K., de la Fuente, G., Harris, P.A., Girdwood, S.E., Pinloche, E., Geor, R.J., Nielsen,
 571 B.D., Schott, H.C., 2nd, Elzinga, S., Newbold, C.J., 2014. Characterisation of the faecal
 572 bacterial community in adult and elderly horses fed a high fibre, high oil or high starch diet
 573 using 454 pyrosequencing. PLoS One. 9, e87424.
 574 Duarte, A.M., Jenkins, T.P., Latrofa, M.S., Giannelli, A., Papadopoulos, E., de Carvalho,
 575 L.M., Nolan, M.J., Otranto, D., Cantacessi, C., 2016. Helminth infections and gut microbiota
 576 - a feline perspective. Parasit. Vectors. 9, 625.
 577 El Temsahy, M.M., Ibrahim, I.R., Mossallam, S.F., Mahrous, H., Abdel Bary, A., Abdel
 578 Salam, S.A., 2015. Evaluation of newly isolated probiotics in the protection against
 579 experimental intestinal trichinellosis. Vet. Parasitol. 214, 303-314.
 580 Fernandes, K.A., Kittelmann, S., Rogers, C.W., Gee, E.K., Bolwell, C.F., Bermingham, E.N.,
 581 Thomas, D.G., 2014. Faecal microbiota of forage-fed horses in New Zealand and the
 582 population dynamics of microbial communities following dietary change. PLoS One. 9,
 583 e112846.
 584 Fricke, W.F., Song, Y., Wang, A.J., Smith, A., Grinchuk, V., Mongodin, E., Pei, C., Ma, B.,
 585 Lu, N., Urban, J.F., Jr., Shea-Donohue, T., Zhao, A., 2015. Type 2 immunity-dependent
 586 reduction of segmented filamentous bacteria in mice infected with the helminthic parasite
 587 *Nippostrongylus brasiliensis*. Microbiome. 3, 40.

588 Gause, W.C., Maizels, R.M., 2016. Macrobiota - helminths as active participants and partners
589 of the microbiota in host intestinal homeostasis. *Curr. Opin. Microbiol.* 32, 14-18.

590 Giacomini, P., Zakrzewski, M., Croese, J., Su, X., Sotillo, J., McCann, L., Navarro, S.,
591 Mitreva, M., Krause, L., Loukas, A., Cantacessi, C., 2015. Experimental hookworm infection
592 and escalating gluten challenges are associated with increased microbial richness in celiac
593 subjects. *Sci. Rep.* 5, 13797.

594 Giacomini, P., Zakrzewski, M., Jenkins, T.P., Su, X., Al-Hallaf, R., Croese, J., de Vries, S.,
595 Grant, A., Mitreva, M., Loukas, A., Krause, L., Cantacessi, C., 2016. Changes in duodenal
596 tissue-associated microbiota following hookworm infection and consecutive gluten
597 challenges in humans with coeliac disease. *Sci. Rep.* 6, 36797.

598 Glendinning, L., Nausch, N., Free, A., Taylor, D.W., Mutapi, F., 2014. The microbiota and
599 helminths: sharing the same niche in the human host. *Parasitology.* 141, 1255-1271.

600 Gómez, J.M., González-Megías, A., 2007. Long-term effects of ungulates on phytophagous
601 insects. *Ecol. Entomol.* 32, 229-234.

602 He, X., McLean, J.S., Edlund, A., Yooseph, S., Hall, A.P., Liu, S.Y., Dorrestein, P.C.,
603 Esquenazi, E., Hunter, R.C., Cheng, G., Nelson, K.E., Lux, R., Shi, W., 2015. Cultivation of
604 a human-associated TM7 phylotype reveals a reduced genome and epibiotic parasitic
605 lifestyle. *Proc. Natl. Acad. Sci. U S A.* 112, 244-249.

606 Hinney, B., Wirtherle, N.C., Kyule, M., Miethe, N., Zessin, K.H., Clausen, P.H., 2011.
607 Prevalence of helminths in horses in the state of Brandenburg, Germany. *Parasitol. Res.* 108,
608 1083-1091.

609 Holm, J.B., Sorobetea, D., Kiilerich, P., Ramayo-Caldas, Y., Estelle, J., Ma, T., Madsen, L.,
610 Kristiansen, K., Svensson-Frej, M., 2015. Chronic *Trichuris muris* infection decreases
611 diversity of the intestinal microbiota and concomitantly increases the abundance of
612 *Lactobacilli*. *PLoS One.* 10, e0125495.

613 Houlden, A., Hayes, K.S., Bancroft, A.J., Worthington, J.J., Wang, P., Grencis, R.K.,
 614 Roberts, I.S., 2015. Chronic *Trichuris muris* infection in C57BL/6 mice causes significant
 615 changes in host microbiota and metabolome: Effects reversed by pathogen clearance. PLoS
 616 One. 10, e0125945.

617 Hutchings, M.R., Kyriazakis, I., Papachristou, T.G., Gordon, I.J., Jackson, F., 2000. The
 618 herbivores' dilemma: trade-offs between nutrition and parasitism in foraging decisions.
 619 Oecologia. 124, 242-251.

620 Jensen, A.N., Mejer, H., Molbak, L., Langkjaer, M., Jensen, T.K., Angen, O., Martinussen,
 621 T., Klitgaard, K., Baggesen, D.L., Thamsborg, S.M., Roepstorff, A., 2011. The effect of a
 622 diet with fructan-rich chicory roots on intestinal helminths and microbiota with special focus
 623 on Bifidobacteria and Campylobacter in piglets around weaning. Animal. 5, 851-860.

624 Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., Glockner, F.O.,
 625 2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-
 626 generation sequencing-based diversity studies. Nucleic Acids Res. 41, e1.

627 Kreisinger, J., Bastien, G., Hauffe, H.C., Marchesi, J., Perkins, S.E., 2015. Interactions
 628 between multiple helminths and the gut microbiota in wild rodents. Philos. Trans. R. Soc.
 629 Lond. B. Biol. Sci. 370.

630 Kuehbachner, T., Rehman, A., Lepage, P., Hellmig, S., Folsch, U.R., Schreiber, S., Ott, S.J.,
 631 2008. Intestinal TM7 bacterial phylogenies in active inflammatory bowel disease. J. Med.
 632 Microbiol. 57, 1569-1576.

633 Lee, S.C., Tang, M.S., Lim, Y.A., Choy, S.H., Kurtz, Z.D., Cox, L.M., Gundra, U.M., Cho,
 634 I., Bonneau, R., Blaser, M.J., Chua, K.H., Loke, P., 2014. Helminth colonization is associated
 635 with increased diversity of the gut microbiota. PLoS Negl. Trop. Dis. 8, e2880.

636 Leung, J.M., Loke, P., 2013. A role for IL-22 in the relationship between intestinal helminths,
 637 gut microbiota and mucosal immunity. Int. J. Parasitol. 43, 253-257.

638 Li, R.W., Li, W., Sun, J., Yu, P., Baldwin, R.L., Urban, J.F., 2016. The effect of helminth
 639 infection on the microbial composition and structure of the caprine abomasal microbiome.
 640 Sci. Rep. 6, 20606.

641 Li, R.W., Wu, S., Li, C.J., Li, W., Schroeder, S.G., 2015. Splice variants and regulatory
 642 networks associated with host resistance to the intestinal worm *Cooperia oncophora* in cattle.
 643 Vet. Parasitol. 211, 241-50.

644 Li, R.W., Wu, S., Li, W., Huang, Y., Gasbarre, L.C., 2011. Metagenome plasticity of the
 645 bovine abomasal microbiota in immune animals in response to *Ostertagia ostertagi* infection.
 646 PLoS One. 6, e24417.

647 Li, R.W., Wu, S., Li, W., Navarro, K., Couch, R.D., Hill, D., Urban, J.F., Jr., 2012.
 648 Alterations in the porcine colon microbiota induced by the gastrointestinal nematode
 649 *Trichuris suis*. Infect. Immun. 80, 2150-2157.

650 Lwin, K.O., Matsui, H., 2014. Comparative analysis of the methanogen diversity in horse and
 651 pony by using mcrA gene and archaeal 16s rRNA gene clone libraries. Archaea. 2014,
 652 483574.

653 Lyons, E.T., Drudge, J.H., Tolliver, S.C., 2000. Larval cyathostomiasis. Vet. Clin. North.
 654 Am. Equine Pract. 16, 501-513.

655 Love, S., Murphy, D., Mellor, D., 1999. Pathogenicity of cyathostome infection. Vet.
 656 Parasitol. 85, 113-121.

657 Manichanh, C., Rigottier-Gois, L., Bonnaud, E., Gloux, K., Pelletier, E., Frangeul, L., Nalin,
 658 R., Jarrin, C., Chardon, P., Marteau, P., Roca, J., Dore, J., 2006. Reduced diversity of faecal
 659 microbiota in Crohn's disease revealed by a metagenomic approach. Gut. 55, 205-211.

660 Martinez-Gomez, F., Fuentes-Castro, B.E., Bautista-Garfías, C.R., 2011. The intraperitoneal
 661 inoculation of *Lactobacillus casei* in mice induces total protection against *Trichinella spiralis*
 662 infection at low challenge doses. Parasitol. Res. 109, 1609-1617.

663 Martinez-Gomez, F., Santiago-Rosales, R., Ramon Bautista-Garfias, C., 2009. Effect of
 664 *Lactobacillus casei* Shirota strain intraperitoneal administration in CD1 mice on the
 665 establishment of *Trichinella spiralis* adult worms and on IgA anti-T. spiralis production. Vet.
 666 Parasitol. 162, 171-175.

667 Maruo, T., Sakamoto, M., Ito, C., Toda, T., Benno, Y., 2008. *Adlercreutzia equolifaciens*
 668 gen. nov., sp. nov., an equol-producing bacterium isolated from human faeces, and emended
 669 description of the genus *Eggerthella*. Int. J. Syst. Evol. Microbiol. 58, 1221-1227.

670 Matthews, J.B., 2011. Facing the threat of equine parasitic disease. Equine. Vet. J. 43, 126-
 671 132.

672 McAllister, T.A., Okine, E.K., Mathison, G.W., Cheng, K.-J., 1996. Dietary, environmental
 673 and microbiological aspects of methane production in ruminants. Can. J. Anim. Sci. 76, 231-
 674 243.

675 McKenney, E.A., Williamson, L., Yoder, A.D., Rawls, J.F., Bilbo, S.D., Parker, W., 2015.
 676 Alteration of the rat cecal microbiome during colonization with the helminth *Hymenolepis*
 677 *diminuta*. Gut Microbes. 6, 182-193.

678 Mfitilodze, M.W., Hutchinson, G.W., 1990. Prevalence and abundance of equine strongyles
 679 (Nematoda: Strongyloidea) in tropical Australia. J. Parasitol. 76, 487-494.

680 Mikaelyan, A., Thompson, C.L., Meuser, K., Zheng, H., Rani, P., Plarre, R., Brune, A., 2017.
 681 High-resolution phylogenetic analysis of Endomicrobia reveals multiple acquisitions of
 682 endosymbiotic lineages by termite gut flagellates. Environ. Microbiol. Rep.

683 Morariu, S., Mederle, N., Badea, C., Darabus, G., Ferrari, N., Genchi, C., 2016. The
 684 prevalence, abundance and distribution of cyathostomins (small strongyles) in horses from
 685 Western Romania. Vet. Parasitol. 223, 205-209.

686 Murphy, D., Love, S., 1997. The pathogenic effects of experimental cyathostome infections
 687 in ponies. Vet. Parasitol. 70, 99-110

688 Nielsen, M.K., Baptiste, K.E., Tolliver, S.C., Collins, S.S., Lyons, E.T., 2010. Analysis of
 689 multiyear studies in horses in Kentucky to ascertain whether counts of eggs and larvae per
 690 gram of feces are reliable indicators of numbers of strongyles and ascarids present. Vet.
 691 Parasitol. 174, 77-84.

692 O'Donnell, M.M., Harris, H.M., Jeffery, I.B., Claesson, M.J., Young, B., PW, O.T., Ross,
 693 R.P., 2013. The core faecal bacterial microbiome of Irish Thoroughbred racehorses. Lett.
 694 Appl. Microbiol. 57, 492-501.

695 Nielsen, M.K., Reinemeyer, C.R., Donecker, J.M., Leathwick, D.M., Marchiondo, A.A.,
 696 Kaplan, R.M., 2014. Anthelmintic resistance in equine parasites--current evidence and
 697 knowledge gaps. Vet. Parasitol. 204, 55-63.

698 Oliveira-Sequeira, T.C., David, E.B., Ribeiro, C., Guimaraes, S., Masseno, A.P., Katagiri, S.,
 699 Sequeira, J.L., 2014. Effect of *Bifidobacterium animalis* on mice infected with *Strongyloides*
 700 *venezuelensis*. Rev. Inst. Med. Trop. Sao Paulo 56, 105-109.

701 Peachey, L.E., Jenkins, T.P., Cantacessi, C., 2017. This gut ain't big enough for both of us.
 702 Or is it? Helminth-microbiota interactions in veterinary species. Trends Parasitol. 33, 619-
 703 632.

704 Peregrine, A.S., McEwen, B., Bienzle, D., Koch, T.G., Weese, J.S., 2006. Larval
 705 cyathostomiasis in horses in Ontario: An emerging disease? Can. Vet. J. 47, 80-82.

706 Peregrine, A.S., Molento, M.B., Kaplan, R.M., Nielsen, M.K., 2014. Anthelmintic resistance
 707 in important parasites of horses: does it really matter? Vet. Parasitol. 201, 1-8.

708 Petkevicius, S., Bach Knudsen, K.E., Murrell, K.D., Wachmann, H., 2003. The effect of
 709 inulin and sugar beet fibre on *Oesophagostomum dentatum* infection in pigs. Parasitology.
 710 127, 61-68.

711 Petkevicius, S., Murrell, K.D., Bach Knudsen, K.E., Jorgensen, H., Roepstorff, A., Laue, A.,
 712 Wachmann, H., 2004. Effects of short-chain fatty acids and lactic acids on survival of
 713 *Oesophagostomum dentatum* in pigs. Vet. Parasitol. 122, 293-301.
 714 Petkevicius, S., Thomsen, L.E., Bach Knudsen, K.E., Murrell, K.D., Roepstorff, A., Boes, J.,
 715 2007. The effect of inulin on new and on patent infections of *Trichuris suis* in growing pigs.
 716 Parasitology. 134, 121-127.
 717 Plieskatt, J.L., Deenonpoe, R., Mulvenna, J.P., Krause, L., Sripa, B., Bethony, J.M., Brindley,
 718 P.J., 2013. Infection with the carcinogenic liver fluke *Opisthorchis viverrini* modifies
 719 intestinal and biliary microbiome. FASEB J. 27, 4572-4584.
 720 Proudman, C.J., Hunter, J.O., Darby, A.C., Escalona, E.E., Batty, C., Turner, C., 2015.
 721 Characterisation of the faecal metabolome and microbiome of Thoroughbred racehorses.
 722 Equine. Vet. J. 47, 580-586.
 723 Rausch, S., Held, J., Fischer, A., Heimesaat, M.M., Kuhl, A.A., Bereswill, S., Hartmann, S.,
 724 2013. Small intestinal nematode infection of mice is associated with increased enterobacterial
 725 loads alongside the intestinal tract. PLoS One. 8, e74026.
 726 Rehbein, S., Lindner, T., Visser, M., Winter, R., 2011. Evaluation of a double centrifugation
 727 technique for the detection of *Anoplocephala* eggs in horse faeces. J. Helminthol. 85, 409-
 728 414.
 729 Relf, V.E., Lester, H.E., Morgan, E.R., Hodgkinson, J.E., Matthews, J.B., 2014. Int. J.
 730 Parasitol. 44, 507-514.
 731 Reynolds, L.A., Smith, K.A., Filbey, K.J., Harcus, Y., Hewitson, J.P., Redpath, S.A., Valdez,
 732 Y., Yebra, M.J., Finlay, B.B., Maizels, R.M., 2014a. Commensal-pathogen interactions in the
 733 intestinal tract: *lactobacilli* promote infection with, and are promoted by, helminth parasites.
 734 Gut Microbes. 5, 522-532.

735 Reynolds, L.A., Harcus, Y., Smith, K.A., Webb, L.M., Hewitson, J.P., Ross, E.A., Brown, S.,
 736 Uematsu, S., Akira, S., Gray, D., Gray, M., MacDonald, A.S., Cunningham, A.F., Maizels,
 737 R.M., 2014b. MyD88 signaling inhibits protective immunity to the gastrointestinal helminth
 738 parasite *Heligmosomoides polygyrus*. J. Immunol. 193, 2984-2993.
 739 Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S., 2011.
 740 Metagenomic biomarker discovery and explanation. Genome Biol. 12.
 741 Sepehri, S., Kotlowski, R., Bernstein, C.N., Krause, D.O., 2007. Microbial diversity of
 742 inflamed and noninflamed gut biopsy tissues in inflammatory bowel disease. Inflamm. Bowel
 743 Dis. 13, 675-683.
 744 Shepherd, M.L., Swecker, W.S., Jr., Jensen, R.V., Ponder, M.A., 2012. Characterization of
 745 the faecal bacteria communities of forage-fed horses by pyrosequencing of 16S rRNA V4
 746 gene amplicons. FEMS Microbiol. Lett. 326, 62-68.
 747 Shin, N.R., Whon, T.W., Bae, J.W., 2015. Proteobacteria: microbial signature of dysbiosis in
 748 gut microbiota. Trends Biotechnol. 33, 496-503.
 749 Slapeta, J., Dowd, S.E., Alanazi, A.D., Westman, M.E., Brown, G.K., 2015. Differences in
 750 the faecal microbiome of non-diarrhoeic clinically healthy dogs and cats associated with
 751 *Giardia duodenalis* infection: impact of hookworms and coccidia. Int. J. Parasitol. 45, 585-
 752 594.
 753 St-Pierre, B., Cersosimo, L.M., Ishaq, S.L., Wright, A.D., 2015. Toward the identification of
 754 methanogenic archaeal groups as targets of methane mitigation in livestock animalsr. Front.
 755 Microbiol. 6, 776.
 756 Steelman, S.M., Chowdhary, B.P., Dowd, S., Suchodolski, J., Janecka, J.E., 2012.
 757 Pyrosequencing of 16S rRNA genes in fecal samples reveals high diversity of hindgut
 758 microflora in horses and potential links to chronic laminitis. BMC Vet. Res. 8, 231.

759 Stratford, C.H., McGorum, B.C., Pickles, K.J., Matthews, J.B., 2011. An update on
760 cyathostomins: anthelmintic resistance and diagnostic tools. *Equine Vet. J. Suppl.*, 133-139.

761 Thomsen, L.E., Petkevicius, S., Bach Knudsen, K.E., Roepstorff, A., 2005. The influence of
762 dietary carbohydrates on experimental infection with *Trichuris suis* in pigs. *Parasitology*.
763 131, 857-865.

764 Traversa, D., Iorio, R., Klei, T.R., Kharchenko, V.A., Gawor, J., Otranto, D., Sparagano,
765 O.A., 2007. New method for simultaneous species-specific identification of equine strongyles
766 (nematoda, strongylida) by reverse line blot hybridization. *J. Clin. Microbiol.* 45, 2937-2942.

767 Tuomisto, H., 2010. A consistent terminology for quantifying species diversity? Yes, it does
768 exist. *Oecologia*. 164, 853-860.

769 Uhlinger, C.A., 1991. Equine Small Strongyles - Epidemiology, Pathology, and Control.
770 *Comp. Cont. Educ. Pract.* 13, 863-869.

771 van Klink, R., van der Plas, F., van Noordwijk, C.G., WallisDeVries, M.F., Olff, H., 2015.
772 Effects of large herbivores on grassland arthropod diversity. *Biol. Rev. Camb. Philos. Soc.*
773 90, 347-366.

774 Walk, S.T., Blum, A.M., Ewing, S.A., Weinstock, J.V., Young, V.B., 2010. Alteration of the
775 murine gut microbiota during infection with the parasitic helminth *Heligmosomoides*
776 *polygyrus*. *Inflamm. Bowel Dis.* 16, 1841-1849.

777 Weese, J.S., Holcombe, S.J., Embertson, R.M., Kurtz, K.A., Roessner, H.A., Jalali, M.,
778 Wismer, S.E., 2014. Changes in the faecal microbiota of mares precede the development of
779 post partum colic. *Equine Vet. J.* 47, 641-9.

780 Winker, S., Woese, C.R., 1991. A definition of the domains Archaea, Bacteria and Eucarya in
781 terms of small subunit ribosomal RNA characteristics. *Syst. Appl. Microbiol.* 14, 305-310.

782 Woese, C.R., Gupta, R., 1981. Are archaebacteria merely derived 'prokaryotes'? *Nature*. 289,
783 95-96.

784 Wu, S., Li, R.W., Li, W., Beshah, E., Dawson, H.D., Urban, J.F., Jr., 2012. Worm burden-
785 dependent disruption of the porcine colon microbiota by *Trichuris suis* infection. PLoS One.
786 7, e35470.
787
788
789

790 **Table 1: Faecal egg counts (FEC) recorded from *Chigh* (FEC >100 eggs per gram,**
791 **e.p.g.) and *Clow* (FEC <100 e.p.g.) broodmares enrolled in the study, as well as of non-**
792 **pregnant controls (NPC), over three consecutive samplings performed pre-anthelmintic**
793 **treatment [D0], as well as at 2 and 14 days post-treatment [D2, D14]. Horses with FEC**
794 **of > 200 e.p.g. (C200) and 0 e.p.g. (C0), are indicated in bold.**

<i>Group</i>	<i>Animal i.d.</i>	<i>Age (yrs)</i>	<i>Tapeworm FEC (e.p.g.)</i>	<i>Ascarid FEC (e.p.g.)</i>	<i>Mean consecutive strongyle FEC (e.p.g.) (±SE) D0</i>	<i>FEC D2 (e.p.g.)</i>	<i>FEC D14 (e.p.g.)</i>
<i>Chigh</i>	MA	6	0	0	123 (±19)	0	0
	CT	5	0	0	130 (±14)	0	0
	LE	10	0	0	113 (±17)	0	0
	PB	5	0	0	171 (±15)	0	0
	SC	4	0	0	101 (±8)	2 (±2)	0
	MSJ	6	0	0	100 (±6)	0	0
	WD	7	0	0	128 (±16)	0	0
	NS	8	0	0	120 (±11)	1 (±1)	0
	HY	7	0	0	150 (±14)	0	0
	RM	4	0	0	139 (±49)	0	0
	HS	6	0	0	200 (±39)	1 (±1)	1 (±1)
	HT	6	0	0	228 (±22)	0	0
	NO	6	0	0	271 (±17)	0	1 (±1)
	QM	4	0	0	418 (±112)	0	0
	SB	8	0	0	206 (±37)	1 (±1)	1 (±1)
	TC	4	0	0	235 (±12)	0	1 (±1)
	MQ	6	0	0	228 (±22)	0	0
	VR	4	0	0	279 (±61)	1 (±1)	0
<i>Clow</i>	GL	4	0	0	10 (±2)	0	0
	DWD	8	0	0	0.3 (± 0.3)	0	0
	LVN	4	0	0	5 (±1)	0	0
	BB	7	0	0	1 (±1)	0	4 (±1)
	LF	5	0	0	10 (±2)	0	0
	IR	7	0	0	3 (±1)	0	0
	DDR	8	0	0	1 (±1)	0	0
	LAL	5	0	0	0.3 (±0.3)	0	0
	PT	7	0	0	0.3 (±0.3)	0	0
	PP	6	0	0	2 (±1)	0	0
	MG	5	0	0	0.5 (±0.3)	0	0
	SY	8	0	0	0	0	0
	BX	16	0	0	0	0	0
	MR	10	0	0	0	0	0
	DD		0	0	0	0	0
	ED	8	0	0	0	0	0
	EP	5	0	0	0	0	0
	SWC	8	0	0	0	0	0
NPC	BET	8	0	0	412 (±45)	-	-

795

BL	12	0	0	22 (± 1)	-	-
ST	10	0	0	0	-	-

**The relationships between faecal egg counts and gut microbial composition in UK
Thoroughbreds infected by cyathostomins**

Peachey L.E.¹, Molena R.A.¹, Jenkins T.P.¹, Di Cesare, A.², Traversa, D.², Hodgkinson J.E.³,
Cantacessi C¹.

¹ Department of Veterinary Medicine, University of Cambridge, Madingley Road, CB3 0ES,
United Kingdom

² Faculty of Veterinary Medicine, University of Teramo, Teramo, 64100, Italy

³ Department of Infection Biology, University of Liverpool, Leahurst, Neston, CH64 7TE,
United Kingdom

Corresponding author: Peachey L.E. lep41@cam.ac.uk; Cantacessi C. cc779@cam.ac.uk

Abstract

A growing body of evidence, particularly in humans and rodents, supports the existence of a complex network of interactions occurring between gastrointestinal (GI) helminth parasites and the gut commensal bacteria, with substantial effects on both host immunity and metabolic potential. However, little is known of the fundamental biology of such interactions in other animal species. Nonetheless, given the considerable economic losses associated with GI parasites, particularly in livestock and equines, as well as the global threat of emerging anthelmintic resistance, further explorations of the complexities of host-helminth-microbiota interactions in these species are needed. This study characterises, for the first time, the composition of the equine bacterial commensal flora associated with the presence, in faecal samples, of low (*Clow*) - and high (*Chigh*) numbers of eggs of an important group of equine GI parasites (i.e. the cyathostomins), prior to and following anthelmintic treatment. High-throughput sequencing of microbial 16S rRNA amplicons and associated bioinformatics and statistical analyses of sequence data revealed strong clustering according to faecal egg counts (FEC) ($P = 0.003$). A trend towards increased populations of Methanomicrobia (class) and *Dehalobacterium* (genus) was observed in *Clow* in comparison to *Chigh*. Anthelmintic treatment in *Chigh* was associated with a significant reduction of the bacterial phylum TM7 14 days post-ivermectin administration, as well as a transient expansion of *Adlercreuzia* spp. at 2 days post-treatment. This study provides a first insight into the discovery of the intimate mechanisms governing host-parasite-microbiota interactions in equids, and sets a basis for the development of novel, biology-based intervention strategies against equine GI helminths based on the manipulation of the commensal gut flora.

Key words: Helminth-microbiota interactions, Thoroughbred horses, cyathostomins, 16S rRNA sequencing, *Adlercreuzia*, Methanomicrobia, TM7.

1. Introduction

Cyathostomins are amongst the most important intestinal nematodes of horses globally (Love et al., 1999; Matthews, 2011; Stratford et al., 2011) with reported prevalence rates as high as 89-100% in equine herds (Mfitilodze and Hutchinson, 1990; Collobert-Laugier et al., 2002; Hinney et al., 2011; ~~Mfitilodze and Hutchinson, 1990;~~ Morariu et al., 2016). Clinical signs of cyathostomin infection range from non-specific weight loss to colic and colitis caused by mass emergence of larvae from the large intestinal wall (= larval cyathostominosis), which may prove fatal (Uhlinger, 1991; Murphy and Love, 1997; Lyons et al., 2000; Peregrine et al., 2006). Thoroughbred (TB) young-stock kept in herds are at high risk of developing serious complications of infection, and hence the implementation of effective strategies for parasite control is a top priority for the TB equine industry. ~~Traditionally,~~ eControl of cyathostomin infections ~~relies—has traditionally relied~~ on the regular administration of chemotherapeutics (i.e. anthelmintics); however, the frequent and uncontrolled use of these compounds has led to the global emergence of resistant populations of parasites (Nielsen et al., 2014; Peregrine et al., 2014). In particular, foci of multi-drug resistance have been recently reported in UK TB stud farms (Relf et al., 2014). This observation, coupled with the lack of novel anthelmintic compounds licenced for use in equids, represents a ‘Damocle’s sword’ for the UK (and global) equine industry. Therefore, alternative strategies for parasite control are urgently needed; in order to support the discovery of such strategies, a deeper understanding of the complex interactions occurring at the host-parasite interface, particularly at the site/s of infection (i.e. the gut), is required.

While a multitude of factors are responsible for the host-parasite interactions which determine infection outcome, increasing attention is being paid to the complex interplay between GI parasites and the host commensal gut microbiota (Bancroft et al., 2012; Glendinning et al., 2014). Indeed, recent studies have reported significant fluctuations in the

76 composition of the vertebrate gut microbiota associated with helminth infections, that were
77 accompanied with shifts in both systemic and local immunity (Bancroft et al., 2012; [Leung](#)
78 [and Loke, 2013](#); [Fricke et al., 2015](#); [Houlden et al., 2015](#); Cattadori et al., 2016; ~~Fricke et al.,~~
79 ~~2015~~; Gause and Maizels, 2016; ~~Houlden et al., 2015~~; ~~Leung and Loke, 2013~~). However, thus
80 far, knowledge of helminth–microbiota cross-talk relies heavily on studies conducted in
81 humans and/or rodent models of infection and disease ([Walk et al., 2010](#); [Rausch et al., 2013](#);
82 [Cantacessi et al., 2014](#); ~~Cattadori et al., 2016~~; [Lee et al., 2014](#); [Reynolds et al., 2014a](#); Fricke
83 et al., 2015; [Giacomin et al., 2015](#); ~~Giacomin et al., 2016~~; Holm et al., 2015; Houlden et al.,
84 2015; [Kreisinger et al., 2015](#); [Lee et al., 2014](#); [McKenney et al., 2015](#); ~~Rausch et al., 2013~~;
85 ~~Reynolds et al., 2014b~~; [Walk et al., 2010](#); [Cattadori et al., 2016](#); [Giacomin et al., 2016](#)). ~~In~~
86 ~~particular~~ Indeed, while only ~~a few handful of~~ studies have characterised the
87 composition of the gut microbiota of veterinary species infected by GI helminths [thus far](#)
88 (~~Duarte et al., 2016~~; [Li et al., 2011](#); [Li et al., 2012](#); ~~Li et al., 2016~~; [Slapeta et al., 2015](#); [Wu et](#)
89 [al., 2012](#); ~~Slapeta et al., 2015~~; [Duarte et al., 2016](#); [Li et al., 2016](#); reviewed by [Peachey et al.,](#)
90 2017). ~~no data is thus currently far~~ available on the effects of infections by GI helminths,
91 such as cyathostomins, on the composition of the equine commensal flora. Aquiring this
92 fundamental knowledge will be key to the development of novel holistic approaches to equid
93 parasite control, aimed at improving host reponses to infections. In this study, we
94 characterise, for the first time, the gut microbial profiles of a cohort of UK TB broodmares
95 with low- and high-numbers of cyathostomin eggs in faeces (as determined by faecal egg
96 counts analysis [FEC]), and examine the effects that administration of a commonly used
97 anthelmintic, i.e. ivermectin, exerts on the overall composition of the gut microbiota as well
98 as relative abundances of individual microbial species.

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Not Highlight

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, Highlight

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

2. Materials and Methods

2.1. Ethics statement

This study was approved and carried out in strict accordance and compliance with the guidelines of the Institutional Ethical Review Committee, Department of Veterinary Medicine, University of Cambridge, UK (Research Project No. CR190). Written informed consent was obtained from the stud farm from which study samples were collected.

2.2. Sampling and diagnostic procedures

For this study, a cohort of TB broodmares was recruited from a stud farm in the East of England, UK. The stud hosts a group of ~130 pregnant broodmares each year, which are kept at pasture in groups of 2-8 across a 1200-acre farm. All broodmares are subjected to targeted anthelmintic treatments (ivermectin and moxidectin), based on FEC measurements at three monthly intervals. In addition, praziquantel is administered to each broodmare three times a year for tapeworm control, whilst a single moxidectin treatment is administered in late November for control of encysted cyathostomin larvae. Samples used in this study were collected in September-October 2016; all broodmares had received ivermectin and praziquantel in May and August 2016, respectively. A total of 117 TB pregnant broodmares, between 5 and 8 months of gestation at the time of sampling, were screened for infections by cyathostomins. Briefly, individual faecal samples were collected on three consecutive days over a seven-day period; aliquots of each sample were subjected to FEC analysis using a centrifugal floatation technique sensitive to one egg per gram (e.p.g.) (Christie and Jackson, 1982), and to screening for infections by the common equine cestode *Anoplocephala perfoliata* using a double sugar flotation technique (Rehbein et al., 2011). Upon observation of strongyle eggs during FEC analysis, the remaining faecal aliquots were subjected to larval

culture to allow for subsequent identification of infecting nematode species using an established Reverse Line Blot (RLB) hybridisation method (Traversa et al., 2007; Cwiklinski et al., 2012). Briefly, genomic DNA was extracted from individual third stage larvae (L3s) harvested from each larval culture, and the intergenic spacer (IGS) region was amplified by nested PCR using conserved biotin labelled primers (Traversa et al., 2007). The PCR products were then incubated with biodyne C membrane bound specific DNA probes for 21 different cyathostomin species (Cwiklinski et al., 2012), incubated with extravidin peroxidase and visualised using x-ray film. Horses were recruited in our study if they satisfied the following criteria: (i) FEC of >100 e.p.g. (= *Chigh*) or <10 e.p.g. (*Clow*) in three consecutive samples collected over a seven-day period; (ii) matched by approximate age and paddock; (iii) negative for co-infections with other gastrointestinal (GI) helminths; (iv) no antibiotic treatment for at least two months prior to sampling; and (v) no previous anthelmintic treatment other than praziquantel for at least four months prior to sampling. Horses enrolled were kept at pasture for the duration of the study and fed 1kg of custom concentrate mix daily.

2.3. Anthelmintic treatment

Individual, naturally voided, faecal samples were collected from the centre of the faecal mass from *Chigh* and *Clow* animals, as well as from three non-pregnant broodmares on day 0 (D0). Then, an anthelmintic treatment (Eqvalan: ivermectin 0.2 mg/kg) was immediately administered to each animal. Sampling was repeated as above at day 2 (D2) and day 14 (D14) post-treatment (p.t.). ~~AA~~ 100g aliquot of each faecal sample was snap frozen, transported to the laboratory and stored at -20 °C within 2h of collection, prior to genomic DNA extraction and sequencing of the bacterial 16S rRNA gene (see below), while the remainder was kept fresh. ~~The remainder of each sample was and~~ subjected to FEC analysis as described above.

150

151 *2.4. High-throughput 16S rRNA sequencing*

152 Genomic DNA was extracted from ~~each each-individual~~ faecal sample, as well as from five
153 negative 'blank' (= no DNA) controls, using the PowerSoil® DNA Isolation Kit (Qiagen,
154 Carlsbad, CA, USA), according to the manufacturers' instructions. High-throughput
155 sequencing of the V3-V4 hypervariable region of the bacterial 16S rRNA gene was
156 performed on an Illumina MiSeq platform according to the manufacturers' protocols with
157 minor adjustments. Briefly, the V3-V4 region was PCR amplified using universal primers
158 (Forward, 5'-TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG CCT ACG GGN
159 GGC WGC AG-3'; Reverse, 5'-GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG
160 ACA GGA CTA CHV GGG TAT CTA ATC C-3') (Klindworth et al., 2013), that contained
161 the Illumina adapter over- hang nucleotide sequences, using the NEBNext® Q5® Hot Start
162 HiFi DNA polymerase (New England Biolabs® Inc, Massachusetts, USA) and the following
163 thermocycling protocol: 98 °C/2 min, followed by 20 cycles of 98 °C/15 s, 63 °C/30 s, and
164 72 °C/30 s, and a final elongation step of 72 °C/5 min. Amplicons were purified using
165 AMPure XP PCR Purification beads (Beckman Coulter, Brea, California, USA), and the
166 NEBNext hot start high-fidelity DNA polymerase was used for the index PCR with Nextera
167 XT index primers (Illumina, San Diego, California, USA) according to the following
168 thermocycling protocol: 98 °C/30 s, 8 cycles of 98 °C/10 s, 65 °C/75 s and at 65 °C/5 min.
169 The indexed samples were purified using AMPure XP beads, quantified using the Qubit
170 Quant-iT™ dsDNA Broad-Range Assay Kit (Life Technologies, Carlsbad, California,
171 USA), and equal quantities from each sample pooled. The resulting pooled library was
172 quantified using the NEBNext® Library Quant Kit for Illumina® (New England Biolabs®
173 Inc) and sequenced on an Illumina MiSeq platform using the v3 chemistry (301 bp paired-
174 end reads). ~~Raw sequence data have been deposited in the NCBI Sequence Read Archive~~

~~database under accession number XXXXXX [The accession number will be inserted in the revised version of the manuscript].~~

2.5. Bioinformatics analyses

Raw paired-end Illumina reads were trimmed for 16S rRNA gene primer sequences using Cutadapt (<https://cutadapt.readthedocs.org/en/stable/>), and reads were joined using the Quantitative Insights Into Microbial Ecology (QIIME) software suite (version 1.9.0) (Caporaso et al., 2010). Successfully joined sequences were quality filtered in QIIME using the 'usearch_qf' script with default settings. Then, sequences were clustered into Operational Taxonomic Units (OTUs) on the basis of similarity to known bacterial sequences available in the Greengenes database (v13.8; <http://greengenes.secondgenome.com/>; 97% sequence similarity cut-off) using the UCLUST software; sequences that could not be matched to references in the Greengenes database were clustered *de novo* based on pair-wise sequence identity (97% sequence similarity cut-off). The first selected cluster seed was considered as the representative sequence of each OTU. Then, representative sequences were assigned to taxonomy using the UCLUST software. OTUs assigned to sequences obtained from the no-DNA control samples, as well as singleton OTUs, were removed prior to downstream analysis. Normalisation was carried out by generating a subsampled OTU table by random sampling (without replacement) of the input OTU table using an implementation of the Mersenne twister algorithm (<http://www.numpy.org>). Cumulative-sum scaling (CSS) and log2 transformation were applied to account for the non-normal distribution of taxonomic counts data. Statistical analyses were executed using the Calypso software (cgenome.net/calypso/); samples were clustered using supervised Canonical Correlation Analysis including FEC (*Chigh:Clow*) and time-point (D0, D2 and D14 p.t.) as explanatory variables. Differences in bacterial alpha diversity (Shannon diversity) between groups were

Field Code Changed

Field Code Changed

Field Code Changed

200 evaluated using paired t-test or ANOVA, depending on number of groups for comparison.
201 Beta diversity was calculated using weighted UniFrac distances and based on the matrices,
202 differences in beta diversity were calculated using Permutational Analysis of Multivariate
203 Dispersions (PERMDISP) through the 'betadisper' function (Anderson et al., 2006).
204 Differences in the composition of the faecal microbiota between groups were assessed using
205 the LEfSe (Linear discriminant analysis Effect Size) workflow (Segata et al., 2011), by
206 assigning FEC/timepoint 'groupings' as comparison class. All statistical analyses were
207 repeated on a sub-group of horses with FEC >200 e.p.g. (n=8) and 0 e.p.g. (n=7), hereafter
208 referred to as 'C200' and 'C0', respectively.

209

210 3. Results

211 3.1 Prevalence of cyathostomin infection and group selection

212 Of 117 broodmares examined for cyathostomin infection, 36 matched the criteria outlined in
213 section 2.2. and were therefore enrolled in this study. Of these, 18 broodmares had an average
214 FEC of ≥ 100 e.p.g. (range 100 - 418), and were thus enrolled as *Chigh* (Table 1). Amongst
215 these, 8 were defined as C200 based on average FEC of ≥ 200 e.p.g. (Table 1). Conversely,
216 18 broodmares with FEC of ≤ 10 e.p.g. (range 0 - 10) were enrolled as *Clow* and, of these, 7
217 could be defined as C0 (Table 1). While no known effect size estimates for changes in
218 microbiota due to parasitism in horses are currently available, data from previous studies in
219 other host:helminth systems (effect size: 1.5; cf. Giacomini et al., 2015, 2016) provided us
220 with suggest that the n = 18 animals in each *Chigh* and *Clow* provided 84% power to detect
221 changes in gut microbial composition between groups (n=18 in each *Chigh* and *Clow* group)
222 using canonical analyses of principal coordinates.

Formatted: Font: Italic, No underline,
Font color: Auto

Formatted: Font: Italic, No underline,
Font color: Auto

Larval culture followed by species identification *via* RLB revealed infections by the following cyathostomin species: *Cyathostomum* (*Cya.*) *catinatum* (88%), *Cylicostephanus* (*Cys.*) *longibursatus* (71%), *Cylicocyclus* (*Cyc.*) *nassatus* (59%), *Coronocyclus coronatus* (41%), *Cyc. calicatus* (41%), *Cyc. radiatus* (29%), *Cys. goldi* (24%), *Cys. leptostomus* (18%), *Cya. pateratum* (18%), *Cyc. ashworthi* (18%) ~~and~~, *Cyc. insigne* (12%). FEC analysis performed on samples collected at D2 and D14 p.t. showed ~~faecal egg count~~FEC reduction rates (FECR) of > 95% in all treated animals (Table 1).

230

231 3.2 Microbiota profiling

A total of 39,461,550 raw paired-end reads were generated from DNA faecal extracts of *Chigh* and *Clow* broodmares ~~and~~ subjected to further processing. Following primer trimming, joining of paired-end reads, filtering of low-quality sequences and removal of ‘contaminant’ and singleton OTUs, a total of 8,077,490 (mean = 73,423 ± 3,610) high quality sequences were retained for further bioinformatics analysis (not shown). Raw and curated sequence data generated in this study are available from Mendeley Data at doi:10.17632/g7chkjrp8f.1. The rarefaction curves generated following in-silico subtraction of low-quality and contaminant sequences indicated that the vast majority of faecal bacterial communities were represented in the remaining sequence data, thus allowing us to undertake further analyses. These sequences were assigned to 95,286 OTUs and 15 bacterial phyla, respectively. The phyla Bacteroidetes (39.9%) and Firmicutes (34.0%) were predominant in all samples, followed by the phyla Verrucomicrobia (12.0%), Spirochaetes (3.9%), Fibrobacteres (2.4%), Cyanobacteria (1%), Proteobacteria (0.9%), Euryarcheota (0.4%), Tenericutes (0.4%), TM7 (0.3%), Actinobacteria (0.3%), Lentisphaerae (0.3%), Synergistetes (0.2%), WPS-2 (0.2%) and Planctomycetes (0.1%) (Fig. 1), while 3.4% of OTUs could not be assigned to any bacterial group. Predominant sub-taxa were Bacteroidia (class), Bacteroidales (order) and

Formatted: Font: Times New Roman
12 pt, No underline, Font color: Auto

Bacteroidales (family) within the phylum Bacteroidetes, and Clostridia (class), Clostridiales (order) and Ruminococcae (family) within the Firmicutes (Fig. 1). Two samples, LVN1 and HS2, differed markedly in the relative proportions of the two most abundant phyla, Bacteroidetes and Firmicutes, when compared with samples from other broodmares (Fig. 1), likely indicating dysbiosis. Therefore, in order to reduce biases due to these potential ‘outliers’, these samples were excluded from further statistical analyses (Fig. 1).

254

Figure 1: Bar charts depicting the relative abundances of faecal bacterial phyla from broodmares with faecal egg counts (FEC) >100 eggs per gram (e.p.g.) (= *Chigh*) and <10 e.p.g. (= *Clow*), according to sampling time point (i.e. pre-anthelmintic treatment [D0], and 2 and 14 days post-treatment [D2 and D14, respectively], and in non-pregnant controls (NPC). Samples from broodmares with FEC >200 e.p.g. (*C200*) and 0 (*C0*) are shown in red.

261

3.3. Differences in microbial composition between *Chigh* and *Clow*, and pre- and post-anthelmintic treatment

Microbial community profiles of each sample were grouped by hierarchical clustering and ordinated by supervised CCA. ~~Using these methods, a provided~~ significant association ~~was~~ observed between microbial composition and FEC (*Chigh* versus *Clow*) ($P = 0.003$), while clustering according to time point pre- and post- anthelmintic treatment (D0 versus D14) did not reach statistical significance ($P = 0.686$) (Fig. 2a). CCA of *C200* versus *C0* led to a clear separation according to FEC ($P = 0.001$) ~~(Fig. 2b)~~, whilst the effect of anthelmintic treatment remained insignificant ($P = 0.811$) (Fig. 2b).

271

Figure 2: The microbial composition of faecal samples ordered by supervised canonical correspondence analysis (CCA) from broodmares with (a) faecal egg counts (FEC) >100 eggs per gram (e.p.g.) (= *Chigh*) and <10 e.p.g. (= *Clow*), pre-anthelmintic treatment [D0] and at 14 days post-treatment [D14], (b) with FEC >200 e.p.g. (*C200*) and 0 (*C0*) at D0 and D14.

No significant differences in OTU alpha diversity (Shannon Index) were recorded between *Chigh* and *Clow*, or between samples collected at D0, D2 and D14 p.t. (Fig.3 a,b,c). A trend towards increased alpha diversity in *Chigh* versus *Clow* at all time-points was observed ($P = 0.087$) (Fig 3a). This trend was also observed when consistent in *C200* samples were compared to *versus* *C0* at T-D0, despite smaller group sizes ($P = 0.102$) (Fig. 3c). No significant differences in beta diversity, as measured by PERMDISP, were observed between groups (Fig. 4).

Figure 3: Shannon diversity charts comparing faecal microbial alpha diversity of broodmares (a) with faecal egg counts (FEC) >100 eggs per gram (e.p.g.) (= *Chigh*) and <10 e.p.g. (= *Clow*) at all time points (i.e. pre-anthelmintic treatment [D0] and 2 and 14 days post-treatment [D2 and D14, respectively]), (b) *Chigh* and *Clow* at D0 only, (c) with FEC >200 e.p.g. (*C200*) and 0 (*C0*) at D0 only, and (d) *Chigh* and (according to sampling time point (i.e. D0, D2 and D14) and (e) *Clow* at according to sampling time point (i.e. D0, D2 and D14).

Figure 4: Permutational Analysis of Multivariate Dispersions (PERMDISP) plots comparing the faecal microbial beta diversity of broodmares (a) with faecal egg counts (FEC) >100 eggs per gram (e.p.g.) (*Chigh*) and <10 e.p.g. (*Clow*) at all time points (i.e.

pre-anthelmintic treatment [D0] and 2 and 14 days post-treatment [D2 and D14, respectively]), (b) *Chigh* and *Clow* at D0 only, (c) with FEC >200 e.p.g. (C200) and 0 (C0) at D0 only, and (d) *Chigh* and according to sampling time point (i.e. D0, D2 and D14) and (e) *Clow* according to sampling time point (i.e. at D0, D2 and D14).

Differences in abundance of individual taxa at the phylum, class, order, family, genus and species level were detected between *Chigh* and *Clow* samples, as well as between samples collected at D0, and D2 and D14 p.t. (Fig. 5). Samples from *Clow* at D0 (pre-treatment) were characterised by an increased abundance of Methanobacteria (class), *Dehalobacterium* (genus) and unclassified *Dehalobacterium* and *Ruminococcus* (species) compared with samples from *Chigh* (Fig. 5a). The same taxa were increased in C0 compared with C200, with the addition of methanogens of the family Methanocorpusculaceae belonging to order Methanomicrobiales, class Methanobacteria, phylum Euryarchaeota; order Endomicrobiales, phylum Elusimicrobia; Rickettsiales (order, family, genus, species); family Bacteroidaceae, genus BF311 and species RFN20 (Fig. 5b). The taxa GMD14H09 (order, family, genus, species) of the phylum Proteobacteria were increased in samples from C200 compared with C0 (Fig. 5b). Anthelmintic treatment in *Chigh* was accompanied by a decrease in the phylum TM7 at D14, whenas compared with pre-treatment samples (Fig. 5c). Additionally, the taxa *Adlercreutzia* and R445B were increased at D2 and D14, respectively, compared with D0 samples (Fig. 5c). In *Clow*, treatment was also associated with an increase in R445B (family, genus, species) at D14 (Fig. 5d).

Figure 5: Linear discriminant analysis Effect Size (LEfSe) bar charts depicting differences in abundance of individual bacterial taxa at the phylum, class, order, family, genus and species level in faecal samples from broodmares (a) with faecal egg counts

(FEC) >100 eggs per gram (e.p.g.) (*Chigh*) and <10 e.p.g. (*Clow*), (b) with FEC >200 e.p.g. (*C200*) and 0 (*C0*) at D0, and in (c) *Chigh* and (d) *Clow* according to sampling time point (i.e. pre-anthelmintic treatment [D0] and 2 and 14 days post-treatment [D2 and D14, respectively]) ~~and (d) *Clow* according to sampling time point (i.e. D0, D2 and D14).~~

Formatted: Font: Italic, No underline,
Font color: Auto

4. Discussion

This study is the first to report a significant association between numbers of cyathostomin eggs in faecal samples from UK Thoroughbreds and the composition of the ~~equine-host~~ gut microbiota. A particularly significant shift in microbial profiles was observed when the faecal bacterial populations of a group of broodmares with FEC of >200 e.p.g. were compared with those with observed FEC of 0. These data are consistent with observations from published studies in both humans and other veterinary species, including rodent models of infection and disease ([Lee et al., 2014](#); [Holm et al., 2015](#); [Houlden et al., 2015](#); [McKenney et al., 2015](#); [Duarte et al., 2016](#); ~~[Holm et al., 2015](#)~~; ~~[Houlden et al., 2015](#)~~; ~~[Lee et al., 2014](#)~~; [Li et al., 2016](#); ~~[McKenney et al., 2015](#)~~). In addition, the administration of a routinely used anthelmintic (i.e. ivermectin) to both *Chigh* and *Clow* resulted in further progressive changes of the microbial profiling of treated horses. While such changes ~~were statistically insignificant~~ did not reach statistical significance when ~~tested-analysed~~ using a multivariate model, this trend suggests that parasite-associated modifications in gut microbiota may be transient, and dependent on the presence of live infections, a hypothesis which requires thorough testing.

Overall, the bacterial phyla identified in this study were consistent between groups of animals enrolled in this study; this observation differs; which contrasts from results from previous studies that had reported significant variability in faecal microbial profiling between horses, largely related to ~~differences-variations~~ in diet and, age, and presence of underlying diseases

(Costa et al., 2012; Daly et al., 2012; [Steelman et al., 2012](#); [O' Donnell et al., 2013](#); Dougal et al., 2014; Fernandes et al., 2014; ~~[O' Donnell et al., 2013](#)~~; ~~[Steelman et al., 2012](#)~~; Weese et al., 2014). Thus, our ~~observation-finding~~ likely indicates that the impact of such confounding factors ~~linked to, e.g., diet, paddock, gender, age,~~ was successfully minimised by our study design, and that the recorded differences in microbial composition were indeed associated to parasite infections. ~~In contrast to previous studies in equids, b~~Bacteria belonging to the phylum Bacteroidetes were predominant in animals examined in our study; conversely, ~~unlike~~ other investigations ~~that have had~~ reported Firmicutes as being the most prevalent phylum in the horse gut flora (Costa et al., 2012; [Shepherd et al., 2012](#); [Dougal et al., 2014](#); [Fernandes et al., 2014](#); [Weese et al., 2014](#); Costa et al., 2015a; Costa et al., 2015b; ~~[Dougal et al., 2014](#)~~; ~~[Fernandes et al., 2014](#)~~; Proudman et al., 2015; ~~[Shepherd et al., 2012](#)~~; ~~[Weese et al., 2014](#)~~). Dietary differences between horse cohorts enrolled in this and previous studies are likely to be responsible for this discrepancy (cf. Daly et al., 2012; Fernandes et al., 2014).

Overall, a trend towards increased microbial alpha diversity, i.e. the number of different OTUs in each sample ['richness'] and their relative abundance [= 'evenness'] (Tuomisto, 2010), was observed in samples from *Chigh* compared with those from *Clow* at D0 (pre-anthelmintic treatment) and in *C200* *versus* *C0*, although these differences did not reach statistical significance. Nevertheless, this observation is supported by the results of a number of previous studies in other host:helminth systems, in which the establishment of parasite infections was associated with an overall increase in alpha diversity of the gut microbiota (Broadhurst et al., 2012; [Lee et al., 2014](#); Giacomini et al., 2015; Giacomini et al., 2016; ~~[Lee et al., 2014](#)~~). Given that a number ~~many of~~ inflammatory GI and/or systemic diseases are accompanied by a reduced alpha diversity ([Manichanh et al., 2006](#); [Sepehri et al., 2007](#); Abrahamsson et al., 2012, 2014; ~~[Manichanh et al., 2006](#)~~; ~~[Sepehri et al., 2007](#)~~), the increase in

372 GI microbial diversity observed in the presence of helminth infections has been hypothesized
373 to represent one of the possible mechanisms by which parasites suppress host inflammatory
374 responses, thus ensuring their long-term survival in the host gut (Bancroft et al., 2012;
375 Glendinning et al., 2014). Therefore, the trends towards increased alpha diversity observed in
376 the faecal microbiota of horses moderately infected by cyathostomins may also result from an
377 increase in gut homeostasis promoted by the parasite. Future studies evaluating the
378 prevalence and incidence of equine inflammatory diseases (e.g. inflammatory bowel disease
379 and recurrent airway obstruction) in the presence or absence of parasite infections could
380 represent significant first steps in this area of research.

381
382 | In addition to global microbial diversity, significant variations in the abundance of specific
383 bacterial taxa were observed between groups. In particular, a higher abundance of
384 microorganisms belonging to the class Methanomicrobia was observed in *Clow* (D0) when
385 compared to *Chigh*. This difference was exacerbated in *C200* versus *C0*, with further
386 significant increases in methanogens belonging to class Methanomicrobia recorded in *C0*,
387 thus suggesting a negative correlation between methanogen abundance and FEC.
388 Methanomicrobia belong to the phylum Euryarchaeota, kingdom Archaea and are
389 phylogenetically distinct from bacteria and eukaryotes, although they retain the prokaryote
390 16S rRNA gene ([Woese and Gupta, 1981](#); Winker and Woese, 1991; ~~Woese and Gupta,~~
391 ~~1981~~). Particularly in ruminants, the role of the Archaeal methanogens in the digestion of
392 fibre has been well documented (St-Pierre et al., 2015). In equids, little is known of the
393 functional diversity of methanogens; however, consistent with our findings, a recent study
394 reported Methanomicrobiales as being predominant in the horse gut (Lwin and Matsui,
395 2014). The underlying mechanisms by which GI helminths may be promoting a reduction in
396 populations of methanogens are yet unclear. Similarly to hypotheses formulated for other

397 host:helminth systems, cyathostomins may prevent expansion of methanogens directly, e.g.
398 *via* their excretory secretory products, or indirectly *via* parasite-induced changes in mucosal
399 immunity (reviewed by Peachey et al., 2017). Alternatively, a high abundance of
400 methanogens prior to helminth infections may bias host immune responses against
401 cyathostomins, thus resulting in the observed low (or absent) parasite burdens. Interestingly,
402 some methanogens (i.e. *Methanosphaera stadtmanae*) have been shown to regulate Th17
403 responses in mice ([Blais Lecours et al., 2011](#); Bernatchez et al., 2017; ~~Blais Lecours et al.,~~
404 ~~2011~~); in turn, these responses have been linked to the ability of mice to clear experimental
405 infections by *Heligmosomoides polygyrus* (Reynolds et al., 2014~~ba~~). Mechanistic studies ~~;~~
406 ~~e.g.~~ aimed to evaluate the effects of expanding populations of gut methanogens on host
407 mucosal responses and, in turn, GI helminth establishment, may help elucidating these
408 interactions.

409
410 An increased abundance of Methanomicrobia in *C*low and *C*0 may also be ~~linked the result of~~
411 ~~to~~ other environmental factors that are simultaneously responsible for the low FEC observed.
412 An example ~~of such a factor~~ is ~~represented by the horse~~ grazing behaviour; indeed, it is
413 known that ~~certain some~~ individuals within a herd favour less nutritional swards of grass in
414 order to avoid faecal contamination (Hutchings et al., 2000). In turn, ~~as~~ animal faeces often
415 act as fertilisers ~~and~~, as a result, individuals favouring nutritious grass are exposed to higher
416 numbers of infective larvae. Grazing different swards of grass may also impact on dietary
417 fibre levels, and thus on gut methanogen populations, ~~as~~ observed in ruminants (McAllister et
418 al., 1996). In horses, dietary factors have also been associated with changes in abundance of
419 Methanomicrobia; for example, *Methanocorpusculum* archaea were observed at a median of
420 17.7% in horses fed a forage-grain diet, and at a median of 31.9% in horses maintained on
421 pasture (Fernandes et al., 2014). Differences in grazing behaviour between individuals may

also be accountable for the increased abundance of bacteria of the phylum Elusimicrobia in C0 versus C200 as these taxa are primarily a component of termite hind-gut microbiota (Mikaelyan et al., 2017; Gómez and González-Megías, 2007; van Klink et al., 2015; Mikaelyan et al., 2017). Experimental cyathostomin infections of stabled horses may eliminate the effect of grazing behaviour on gut microbial profiles, although ethical concerns may prevent the execution of such a study in the future.

In contrast to uninfected horses, the faecal microbial profiles of C200 were characterised by an increased abundance of GMD14H09, phylum Proteobacteria, class Deltaproteobacteria. Increases in Proteobacteria abundance have repeatedly been reported in association with helminth infections, e.g. in mice infected by *Trichuris muris* and *H. polygyrus*, pigs infected by *T. suis*, and rabbits infected by *Trichostrongylus retortaeformis* (Cattadori et al., 2016; Holm et al., 2015; Li et al., 2012; Holm et al., 2015; Cattadori et al., 2016). Proteobacteria are known increase in the presence of ~~to be enriched with~~ GI inflammation ~~and can lead to sustained microbial dysbiosis~~ (Shin et al., 2015); hence, the expansion of populations of Proteobacteria in the faecal microbiota of horses with higher infection burdens ~~is unsurprising~~ may be indicative of an inflammatory status of the intestinal tract of these horses at the time of sampling.

One of the objectives of this study was to assess the impact of anthelmintic treatment on the faecal microbial profiling of cyathostomin-infected horses. In particular, ivermectin administration to *Chigh* was followed by a significant decrease in populations of the phylum TM7 at D14. Since the relative abundance of TM7 remained ~~were~~ unchanged following ivermectin administration in *Clow*, it is tempting to speculate that a mutualistic association may exist between TM7 and cyathostomins, whereby each promote establishment of the

other, similarly to ~~the mutual relationship that~~ described for Lactobacillaceae and *H. polygyrus* (Reynolds et al., 2014^{ab}). Bacteria belonging to the phylum TM7 are obligate epibionts of *Actinomyces* spp. (He et al., 2015), and are thus uncultivable. While TM7 ~~has~~ not previously been linked to GI helminth infections^s, this phylum of bacteria has been associated with mucosal inflammatory disease in humans (Kuehbachner et al., 2008). Interestingly, ~~isolates from~~ TM7 isolates have been shown to repress the induction of TNF- α production in macrophages infected by *Actinomyces odontolyticus*, thus suggesting a potential immune suppressive activity (He et al., 2015); ~~therefore~~hence, TM7 may promote the establishment of cyathostomins by suppressing effective anti-parasite immune responses. Furthermore, an increase ~~of~~in the taxa *Adlercreutzia* (phylum actinobacteria) and R445B (phylum Lentisphaerae) was observed in *Chigh* at D2 and D14, respectively. The latter was also increased in *Clow* at D14, suggesting that this change was unrelated to cyathostomin removal. Bacteria of the genus *Adlercreutzia* produce the metabolite equol (Maruo et al., 2008), a known anti-inflammatory and vasodilator (Blay et al., 2010). ~~Hence~~Thus, it could be hypothesised that increases in populations of *Adlercreutzia* and its metabolites following ivermectin administration might contribute to the emergence of hypobiotic larval stages of cyathostomins ~~(, which is known to occur post-anthelmintic treatment;)~~ (Lyons et al., 2000), via ~~suppress~~the suppression of~~ing effective~~ mucosal immune responses~~ity~~. This hypothesis requires testing in controlled mechanistic experiments.

FEC are often utilised as proxy of parasite infection burdens; however, several investigations have confuted this practice, as weak correlations have been detected between FEC and parasite burdens in horses with >500 e.p.g. of faeces (Nielsen et al., 2010). While the FEC cut-offs used in this study are indicative of differing infection burdens between groups, any inference on the relationships between number of worms in the horse intestine and gut microbial profiling must be taken with caution. Ethical considerations prevent us from

performing post-mortem total worm counts on experimentally infected horses; nevertheless, in the future, it may be possible to establish unequivocal relationships between cyathostomin infection burdens (including encysted larvae) and gut microbial profiling from samples collected in abattoir.

Clearly, a complex network of host-parasite interactions, as well as environmental factors, contribute to the findings reported in this study, and thus further work is needed to disentangle the causality of these relationships. However, one key question that needs addressing is whether differences in host immunity may be associated with significant changes in gut microbial composition (and *vice versa*) and, if ~~such is the case~~, whether the horse gut microbiota could be manipulated to improve resistance to helminth infection. Indeed, previous investigations in cattle and mice have reported that host genes encoding for antimicrobial proteins are up-regulated in the mucosa of animals ~~which are bred for~~ resistance to helminth infection (D'Elia et al., 2009; Li et al., 2015). In addition, dietary supplementation with both pro- (Bautista-Garfias et al., 1999; Bautista-Garfias et al., 2001; [Martinez-Gomez et al., 2009](#); [Martinez-Gomez et al., 2011](#); [Oliveira-Sequeira et al., 2014](#); El Tamsahy et al., 2015; ~~Martinez-Gomez et al., 2011~~; ~~Martinez-Gomez et al., 2009~~; ~~Oliveira-Sequeira et al., 2014~~) and pre-biotics (~~Jensen et al., 2011~~; Petkevicius et al., 2003; Petkevicius et al., 2004; [Thomsen et al., 2005](#); Petkevicius et al., 2007; [Jensen et al., 2011](#); ~~Thomsen et al., 2005~~), has led to significant reductions in worm burdens in murine and swine helminth infection models, thus indicating that alterations to the bacterial flora in the host GI tract may bias host immune responses against ~~helminth infection~~ parasites. ~~Consequently,~~ Further characterisation of equine host mucosal responses and GI microbiota, in the presence or absence of helminth infection and accompanied by total enumeration of infecting parasites,

is a key area of future research, as it may lead to the identification of microbial factors linked to host susceptibility.

5. Conclusion

Cyathostomin infection in horses was associated with global shifts in faecal microbial composition and diversity, in accordance with previous studies in other host: helminth systems, as well as significant changes in specific populations of gut bacteria. Such changes predominantly involved ‘minor’ phyla, thus suggesting that the equine ‘core gut microbiota’ remains unaltered in the presence of burdens of cyathostomins such as those observed in this study. Our findings also support the hypothesis that selected bacterial taxa, and/or their metabolites, may play roles in biasing the host immune response either for, e.g. TM7, or against, e.g. Methanomicrobia, cyathostomin infection in horses. These data pave the way for future mechanistic studies aimed to identify microbial factors linked to host susceptibility, and to ~~alter~~manipulate the GI microbiota of horses (e.g. via dietary or probiotic interventions), in order to improve resistance to cyathostomins.

513 **Role of the Funding Body**

514 L.E.P. is a Post-Doctoral Research Fellow co-funded by the Horserace Betting Levy Board,
515 the Thoroughbred Breeders' Association and the British European Breeders Fund (Grant no.
516 VET/EPDF/2016-2). T.P.J. is the grateful recipient of a Ph.D. scholarship by the
517 Biotechnology and Biological Sciences Research Council. Research in C.C. laboratory is
518 funded by grants by the Royal Society and the Isaac Newton Trust/ Wellcome Trust/
519 University of Cambridge.

520

521 **Acknowledgements**

522 The authors are very grateful for the cooperation and assistance provided by the management
523 and staff of the Thoroughbred stud farm where this study was performed. Additionally, the
524 authors would like to thank Professor James Wood ([University of Cambridge](#)) for
525 constructive comments.

526

527 **References**

528 Abrahamsson, T.R., Jakobsson, H.E., Andersson, A.F., Bjorksten, B., Engstrand, L.,
529 Jenmalm, M.C., 2012. Low diversity of the gut microbiota in infants with atopic eczema. *J.*
530 *Allergy Clin. Immunol.* 129, 434-440, 440 e431-432.

531 Abrahamsson, T.R., Jakobsson, H.E., Andersson, A.F., Bjorksten, B., Engstrand, L.,
532 Jenmalm, M.C., 2014. Low gut microbiota diversity in early infancy precedes asthma at
533 school age. *Clin. Exp. Allergy.* 44, 842-850.

534 Anderson, M.J., Ellingsen, K.E., McArdle, B.H., 2006. Multivariate dispersion as a measure
535 of beta diversity. *Ecol. Lett.* 9, 683-693.

536 Bancroft, A.J., Hayes, K.S., Grenis, R.K., 2012. Life on the edge: the balance between
537 macrofauna, microflora and host immunity. *Trends Parasitol.* 28, 93-98.

538 Bautista-Garfias, C.R., Ixta, O., Orduna, M., Martinez, F., Aguilar, B., Cortes, A., 1999.
 539 Enhancement of resistance in mice treated with *Lactobacillus casei*: effect on *Trichinella*
 540 *spiralis* infection. Vet. Parasitol. 80, 251-260.
 541 Bautista-Garfias, C.R., Ixta-Rodriguez, O., Martinez-Gomez, F., Lopez, M.G., Aguilar-
 542 Figueroa, B.R., 2001. Effect of viable or dead *Lactobacillus casei* organisms administered
 543 orally to mice on resistance against *Trichinella spiralis* infection. Parasite. 8, S226-228.
 544 Bernatchez, E., Gold, M.J., Langlois, A., Blais-Lecours, P., Boucher, M., Duchaine, C.,
 545 Marsolais, D., McNaghy, K.M., Blanchet, M.R., 2017. *Methanosphaera stadtmanae* induces
 546 a type IV hypersensitivity response in a mouse model of airway inflammation. Physiol. Rep.
 547 5.
 548 Blais Lecours, P., Duchaine, C., Taillefer, M., Tremblay, C., Veillette, M., Cormier, Y.,
 549 Marsolais, D., 2011. Immunogenic properties of archaeal species found in bioaerosols. PLoS
 550 One. 6, e23326.
 551 Blay, M., Espinel, A.E., Delgado, M.A., Baiges, I., Blade, C., Arola, L., Salvado, J., 2010.
 552 Isoflavone effect on gene expression profile and biomarkers of inflammation. J. Pharm.
 553 Biomed. Anal. 51, 382-390.
 554 Broadhurst, M.J., Ardeshtir, A., Kanwar, B., Mirpuri, J., Gundra, U.M., Leung, J.M., Wiens,
 555 K.E., Vujkovic-Cvijin, I., Kim, C.C., Yarovinsky, F., Lerche, N.W., McCune, J.M., Loke, P.,
 556 2012. Therapeutic helminth infection of macaques with idiopathic chronic diarrhea alters the
 557 inflammatory signature and mucosal microbiota of the colon. PLoS Pathog. 8, e1003000.
 558 Cantacessi, C., Giacomini, P., Croese, J., Zakrzewski, M., Sotillo, J., McCann, L., Nolan,
 559 M.J., Mitreva, M., Krause, L., Loukas, A., 2014. Impact of experimental hookworm infection
 560 on the human gut microbiota. J. Infect. Dis. 210, 1431-1434.
 561 Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K.,
 562 Fierer, N., Pena, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T., Knights, D.,

Field Code Changed

563 Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D., Pirrung, M., Reeder,
 564 J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A., Widmann, J., Yatsunenko, T., Zaneveld,
 565 J., Knight, R., 2010. QIIME allows analysis of high-throughput community sequencing data.
 566 Nat. Methods. 7, 335-336.

567 Cattadori, I.M., Sebastian, A., Hao, H., Katani, R., Albert, I., Eilertson, K.E., Kapur, V.,
 568 Pathak, A., Mitchell, S., 2016. Impact of helminth infections and nutritional constraints on
 569 the small intestine microbiota. PLoS One. 11, e0159770.

570 Christie, M., Jackson, F., 1982. Specific identification of strongyle eggs in small samples of
 571 sheep faeces. Res. Vet. Sci. 32, 113-117.

572 Collobert-Laugier, C., Hoste, H., Sevin, C., Dorchies, P., 2002. Prevalence, abundance and
 573 site distribution of equine small strongyles in Normandy, France. Vet. Parasitol. 110, 77-83.

574 Costa, M.C., Arroyo, L.G., Allen-Vercoe, E., Stampfli, H.R., Kim, P.T., Sturgeon, A.,
 575 Weese, J.S., 2012. Comparison of the fecal microbiota of healthy horses and horses with
 576 colitis by high throughput sequencing of the V3-V5 region of the 16S rRNA gene. PLoS One.
 577 7, e41484.

578 Costa, M.C., Silva, G., Ramos, R.V., Staempfli, H.R., Arroyo, L.G., Kim, P., Weese, J.S.,
 579 2015a. Characterization and comparison of the bacterial microbiota in different
 580 gastrointestinal tract compartments in horses. Vet. J. 205, 74-80.

581 Costa, M.C., Stampfli, H.R., Arroyo, L.G., Allen-Vercoe, E., Gomes, R.G., Weese, J.S.,
 582 2015b. Changes in the equine fecal microbiota associated with the use of systemic
 583 antimicrobial drugs. BMC Vet. Res. 11, 19.

584 Cwiklinski, K., Kooyman, F.N., Van Doorn, D.C., Matthews, J.B., Hodgkinson, J.E., 2012.
 585 New insights into sequence variation in the IGS region of 21 cyathostomin species and the
 586 implication for molecular identification. Parasitology. 139, 1063-1073.

587 D'Elia, R., DeSchoolmeester, M.L., Zeef, L.A., Wright, S.H., Pemberton, A.D., Else, K.J.,
 588 2009. Expulsion of *Trichuris muris* is associated with increased expression of angiogenin 4 in
 589 the gut and increased acidity of mucins within the goblet cell. BMC Genomics. 10, 492.
 590 Daly, K., Proudman, C.J., Duncan, S.H., Flint, H.J., Dyer, J., Shirazi-Beechey, S.P., 2012.
 591 Alterations in microbiota and fermentation products in equine large intestine in response to
 592 dietary variation and intestinal disease. Br. J. Nutr. 107, 989-995.
 593 Dougal, K., de la Fuente, G., Harris, P.A., Girdwood, S.E., Pinloche, E., Geor, R.J., Nielsen,
 594 B.D., Schott, H.C., 2nd, Elzinga, S., Newbold, C.J., 2014. Characterisation of the faecal
 595 bacterial community in adult and elderly horses fed a high fibre, high oil or high starch diet
 596 using 454 pyrosequencing. PLoS One. 9, e87424.
 597 Duarte, A.M., Jenkins, T.P., Latrofa, M.S., Giannelli, A., Papadopoulos, E., de Carvalho,
 598 L.M., Nolan, M.J., Otranto, D., Cantacessi, C., 2016. Helminth infections and gut microbiota
 599 - a feline perspective. Parasit. Vectors. 9, 625.
 600 El Temsahy, M.M., Ibrahim, I.R., Mossallam, S.F., Mahrous, H., Abdel Bary, A., Abdel
 601 Salam, S.A., 2015. Evaluation of newly isolated probiotics in the protection against
 602 experimental intestinal trichinellosis. Vet. Parasitol. 214, 303-314.
 603 Fernandes, K.A., Kittelmann, S., Rogers, C.W., Gee, E.K., Bolwell, C.F., Bermingham, E.N.,
 604 Thomas, D.G., 2014. Faecal microbiota of forage-fed horses in New Zealand and the
 605 population dynamics of microbial communities following dietary change. PLoS One. 9,
 606 e112846.
 607 Fricke, W.F., Song, Y., Wang, A.J., Smith, A., Grinchuk, V., Mongodin, E., Pei, C., Ma, B.,
 608 Lu, N., Urban, J.F., Jr., Shea-Donohue, T., Zhao, A., 2015. Type 2 immunity-dependent
 609 reduction of segmented filamentous bacteria in mice infected with the helminthic parasite
 610 *Nippostrongylus brasiliensis*. Microbiome. 3, 40.

611 Gause, W.C., Maizels, R.M., 2016. Macrobiota - helminths as active participants and partners
612 of the microbiota in host intestinal homeostasis. *Curr. Opin. Microbiol.* 32, 14-18.

613 Giacomini, P., Zakrzewski, M., Croese, J., Su, X., Sotillo, J., McCann, L., Navarro, S.,
614 Mitreva, M., Krause, L., Loukas, A., Cantacessi, C., 2015. Experimental hookworm infection
615 and escalating gluten challenges are associated with increased microbial richness in celiac
616 subjects. *Sci. Rep.* 5, 13797.

617 Giacomini, P., Zakrzewski, M., Jenkins, T.P., Su, X., Al-Hallaf, R., Croese, J., de Vries, S.,
618 Grant, A., Mitreva, M., Loukas, A., Krause, L., Cantacessi, C., 2016. Changes in duodenal
619 tissue-associated microbiota following hookworm infection and consecutive gluten
620 challenges in humans with coeliac disease. *Sci. Rep.* 6, 36797.

621 Glendinning, L., Nausch, N., Free, A., Taylor, D.W., Mutapi, F., 2014. The microbiota and
622 helminths: sharing the same niche in the human host. *Parasitology.* 141, 1255-1271.

623 Gómez, J.M., González-Megías, A., 2007. Long-term effects of ungulates on phytophagous
624 insects. *Ecol. Entomol.* 32, 229-234.

625 He, X., McLean, J.S., Edlund, A., Yooseph, S., Hall, A.P., Liu, S.Y., Dorrestein, P.C.,
626 Esquenazi, E., Hunter, R.C., Cheng, G., Nelson, K.E., Lux, R., Shi, W., 2015. Cultivation of
627 a human-associated TM7 phylotype reveals a reduced genome and epibiotic parasitic
628 lifestyle. *Proc. Natl. Acad. Sci. U S A.* 112, 244-249.

629 Hinney, B., Wirtherle, N.C., Kyule, M., Miethe, N., Zessin, K.H., Clausen, P.H., 2011.
630 Prevalence of helminths in horses in the state of Brandenburg, Germany. *Parasitol. Res.* 108,
631 1083-1091.

632 Holm, J.B., Sorobetea, D., Kiilerich, P., Ramayo-Caldas, Y., Estelle, J., Ma, T., Madsen, L.,
633 Kristiansen, K., Svensson-Frej, M., 2015. Chronic *Trichuris muris* infection decreases
634 diversity of the intestinal microbiota and concomitantly increases the abundance of
635 *Lactobacilli*. *PLoS One.* 10, e0125495.

636 Houlden, A., Hayes, K.S., Bancroft, A.J., Worthington, J.J., Wang, P., Grencis, R.K.,
 637 Roberts, I.S., 2015. Chronic *Trichuris muris* infection in C57BL/6 mice causes significant
 638 changes in host microbiota and metabolome: Effects reversed by pathogen clearance. PLoS
 639 One. 10, e0125945.
 640 Hutchings, M.R., Kyriazakis, I., Papachristou, T.G., Gordon, I.J., Jackson, F., 2000. The
 641 herbivores' dilemma: trade-offs between nutrition and parasitism in foraging decisions.
 642 Oecologia. 124, 242-251.
 643 Jensen, A.N., Mejer, H., Molbak, L., Langkjaer, M., Jensen, T.K., Angen, O., Martinussen,
 644 T., Klitgaard, K., Baggesen, D.L., Thamsborg, S.M., Roepstorff, A., 2011. The effect of a
 645 diet with fructan-rich chicory roots on intestinal helminths and microbiota with special focus
 646 on Bifidobacteria and Campylobacter in piglets around weaning. Animal. 5, 851-860.
 647 Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., Glockner, F.O.,
 648 2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-
 649 generation sequencing-based diversity studies. Nucleic Acids Res. 41, e1.
 650 Kreisinger, J., Bastien, G., Hauffe, H.C., Marchesi, J., Perkins, S.E., 2015. Interactions
 651 between multiple helminths and the gut microbiota in wild rodents. Philos. Trans. R. Soc.
 652 Lond. B. Biol. Sci. 370.
 653 Kuehbachner, T., Rehman, A., Lepage, P., Hellmig, S., Folsch, U.R., Schreiber, S., Ott, S.J.,
 654 2008. Intestinal TM7 bacterial phylogenies in active inflammatory bowel disease. J. Med.
 655 Microbiol. 57, 1569-1576.
 656 Lee, S.C., Tang, M.S., Lim, Y.A., Choy, S.H., Kurtz, Z.D., Cox, L.M., Gundra, U.M., Cho,
 657 I., Bonneau, R., Blaser, M.J., Chua, K.H., Loke, P., 2014. Helminth colonization is associated
 658 with increased diversity of the gut microbiota. PLoS Negl. Trop. Dis. 8, e2880.
 659 Leung, J.M., Loke, P., 2013. A role for IL-22 in the relationship between intestinal helminths,
 660 gut microbiota and mucosal immunity. Int. J. Parasitol. 43, 253-257.

661 Li, R.W., Li, W., Sun, J., Yu, P., Baldwin, R.L., Urban, J.F., 2016. The effect of helminth
 662 infection on the microbial composition and structure of the caprine abomasal microbiome.
 663 Sci. Rep. 6, 20606.

664 Li, R.W., Wu, S., Li, C.J., Li, W., Schroeder, S.G., 2015. Splice variants and regulatory
 665 networks associated with host resistance to the intestinal worm *Cooperia oncophora* in cattle.
 666 Vet. Parasitol. 211, 241-50.

667 Li, R.W., Wu, S., Li, W., Huang, Y., Gasbarre, L.C., 2011. Metagenome plasticity of the
 668 bovine abomasal microbiota in immune animals in response to *Ostertagia ostertagi* infection.
 669 PLoS One. 6, e24417.

670 Li, R.W., Wu, S., Li, W., Navarro, K., Couch, R.D., Hill, D., Urban, J.F., Jr., 2012.
 671 Alterations in the porcine colon microbiota induced by the gastrointestinal nematode
 672 *Trichuris suis*. Infect. Immun. 80, 2150-2157.

673 Lwin, K.O., Matsui, H., 2014. Comparative analysis of the methanogen diversity in horse and
 674 pony by using mcrA gene and archaeal 16s rRNA gene clone libraries. Archaea. 2014,
 675 483574.

676 Lyons, E.T., Drudge, J.H., Tolliver, S.C., 2000. Larval cyathostomiasis. Vet. Clin. North.
 677 Am. Equine Pract. 16, 501-513.

678 Love, S., Murphy, D., Mellor, D., 1999. Pathogenicity of cyathostome infection. Vet.
 679 Parasitol. 85, 113-121.

680 Manichanh, C., Rigottier-Gois, L., Bonnaud, E., Gloux, K., Pelletier, E., Frangeul, L., Nalin,
 681 R., Jarrin, C., Chardon, P., Marteau, P., Roca, J., Dore, J., 2006. Reduced diversity of faecal
 682 microbiota in Crohn's disease revealed by a metagenomic approach. Gut. 55, 205-211.

683 Martinez-Gomez, F., Fuentes-Castro, B.E., Bautista-Garfias, C.R., 2011. The intraperitoneal
 684 inoculation of *Lactobacillus casei* in mice induces total protection against *Trichinella spiralis*
 685 infection at low challenge doses. Parasitol. Res. 109, 1609-1617.

686 Martinez-Gomez, F., Santiago-Rosales, R., Ramon Bautista-Garfias, C., 2009. Effect of
 687 *Lactobacillus casei* Shirota strain intraperitoneal administration in CD1 mice on the
 688 establishment of *Trichinella spiralis* adult worms and on IgA anti-T. spiralis production. Vet.
 689 Parasitol. 162, 171-175.

690 Maruo, T., Sakamoto, M., Ito, C., Toda, T., Benno, Y., 2008. *Adlercreutzia equolifaciens*
 691 gen. nov., sp. nov., an equol-producing bacterium isolated from human faeces, and emended
 692 description of the genus *Eggerthella*. Int. J. Syst. Evol. Microbiol. 58, 1221-1227.

693 Matthews, J.B., 2011. Facing the threat of equine parasitic disease. Equine. Vet. J. 43, 126-
 694 132.

695 McAllister, T.A., Okine, E.K., Mathison, G.W., Cheng, K.-J., 1996. Dietary, environmental
 696 and microbiological aspects of methane production in ruminants. Can. J. Anim. Sci. 76, 231-
 697 243.

698 McKenney, E.A., Williamson, L., Yoder, A.D., Rawls, J.F., Bilbo, S.D., Parker, W., 2015.
 699 Alteration of the rat cecal microbiome during colonization with the helminth *Hymenolepis*
 700 *diminuta*. Gut Microbes. 6, 182-193.

701 Mfitilodze, M.W., Hutchinson, G.W., 1990. Prevalence and abundance of equine strongyles
 702 (Nematoda: Strongyloidea) in tropical Australia. J. Parasitol. 76, 487-494.

703 Mikaelyan, A., Thompson, C.L., Meuser, K., Zheng, H., Rani, P., Plarre, R., Brune, A., 2017.
 704 High-resolution phylogenetic analysis of Endomicrobia reveals multiple acquisitions of
 705 endosymbiotic lineages by termite gut flagellates. Environ. Microbiol. Rep.

706 Morariu, S., Mederle, N., Badea, C., Darabus, G., Ferrari, N., Genchi, C., 2016. The
 707 prevalence, abundance and distribution of cyathostomins (small strongyles) in horses from
 708 Western Romania. Vet. Parasitol. 223, 205-209.

709 Murphy, D., Love, S., 1997. The pathogenic effects of experimental cyathostome infections
 710 in ponies. Vet. Parasitol. 70, 99-110

711 Nielsen, M.K., Baptiste, K.E., Tolliver, S.C., Collins, S.S., Lyons, E.T., 2010. Analysis of
712 multiyear studies in horses in Kentucky to ascertain whether counts of eggs and larvae per
713 gram of feces are reliable indicators of numbers of strongyles and ascarids present. Vet.
714 Parasitol. 174, 77-84.

715 O'Donnell, M.M., Harris, H.M., Jeffery, I.B., Claesson, M.J., Younge, B., PW, O.T., Ross,
716 R.P., 2013. The core faecal bacterial microbiome of Irish Thoroughbred racehorses. Lett.
717 Appl. Microbiol. 57, 492-501.

718 ~~Nielsen, M.K., Baptiste, K.E., Tolliver, S.C., Collins, S.S., Lyons, E.T., 2010. Analysis of~~
719 ~~multiyear studies in horses in Kentucky to ascertain whether counts of eggs and larvae per~~
720 ~~gram of feces are reliable indicators of numbers of strongyles and ascarids present. Vet.~~
721 ~~Parasitol. 174, 77-84.~~

722 Nielsen, M.K., Reinemeyer, C.R., Donecker, J.M., Leathwick, D.M., Marchiondo, A.A.,
723 Kaplan, R.M., 2014. Anthelmintic resistance in equine parasites--current evidence and
724 knowledge gaps. Vet. Parasitol. 204, 55-63.

725 Oliveira-Sequeira, T.C., David, E.B., Ribeiro, C., Guimaraes, S., Masseno, A.P., Katagiri, S.,
726 Sequeira, J.L., 2014. Effect of *Bifidobacterium animalis* on mice infected with *Strongyloides*
727 *venezuelensis*. Rev. Inst. Med. Trop. Sao Paulo 56, 105-109.

728 Peachey, L.E., Jenkins, T.P., Cantacessi, C., 2017. This gut ain't big enough for both of us.
729 Or is it? Helminth-microbiota interactions in veterinary species. Trends Parasitol. 33, 619-
730 632.

731 Peregrine, A.S., McEwen, B., Bienzle, D., Koch, T.G., Weese, J.S., 2006. Larval
732 cyathostomiasis in horses in Ontario: An emerging disease? Can. Vet. J. 47, 80-82.

733 Peregrine, A.S., Molento, M.B., Kaplan, R.M., Nielsen, M.K., 2014. Anthelmintic resistance
734 in important parasites of horses: does it really matter? Vet. Parasitol. 201, 1-8.

735 Petkevicius, S., Bach Knudsen, K.E., Murrell, K.D., Wachmann, H., 2003. The effect of
 736 inulin and sugar beet fibre on *Oesophagostomum dentatum* infection in pigs. Parasitology.
 737 127, 61-68.
 738 Petkevicius, S., Murrell, K.D., Bach Knudsen, K.E., Jorgensen, H., Roepstorff, A., Laue, A.,
 739 Wachmann, H., 2004. Effects of short-chain fatty acids and lactic acids on survival of
 740 *Oesophagostomum dentatum* in pigs. Vet. Parasitol. 122, 293-301.
 741 Petkevicius, S., Thomsen, L.E., Bach Knudsen, K.E., Murrell, K.D., Roepstorff, A., Boes, J.,
 742 2007. The effect of inulin on new and on patent infections of *Trichuris suis* in growing pigs.
 743 Parasitology. 134, 121-127.
 744 Plieskatt, J.L., Deenonpoe, R., Mulvenna, J.P., Krause, L., Sripa, B., Bethony, J.M., Brindley,
 745 P.J., 2013. Infection with the carcinogenic liver fluke *Opisthorchis viverrini* modifies
 746 intestinal and biliary microbiome. FASEB J. 27, 4572-4584.
 747 Proudman, C.J., Hunter, J.O., Darby, A.C., Escalona, E.E., Batty, C., Turner, C., 2015.
 748 Characterisation of the faecal metabolome and microbiome of Thoroughbred racehorses.
 749 Equine. Vet. J. 47, 580-586.
 750 Rausch, S., Held, J., Fischer, A., Heimesaat, M.M., Kuhl, A.A., Bereswill, S., Hartmann, S.,
 751 2013. Small intestinal nematode infection of mice is associated with increased enterobacterial
 752 loads alongside the intestinal tract. PLoS One. 8, e74026.
 753 Rehbein, S., Lindner, T., Visser, M., Winter, R., 2011. Evaluation of a double centrifugation
 754 technique for the detection of *Anoplocephala* eggs in horse faeces. J. Helminthol. 85, 409-
 755 414.
 756 Relf, V.E., Lester, H.E., Morgan, E.R., Hodgkinson, J.E., Matthews, J.B., 2014. Int. J.
 757 Parasitol. 44, 507-514.
 758 [Reynolds, L.A., Smith, K.A., Filbey, K.J., Harcus, Y., Hewitson, J.P., Redpath, S.A., Valdez,](#)
 759 [Y., Yebra, M.J., Finlay, B.B., Maizels, R.M., 2014a. Commensal-pathogen interactions in the](#)

760 [intestinal tract: *lactobacilli* promote infection with, and are promoted by, helminth parasites.](#)
761 [Gut Microbes. 5, 522-532.](#)

762 Reynolds, L.A., Harcus, Y., Smith, K.A., Webb, L.M., Hewitson, J.P., Ross, E.A., Brown, S.,
763 Uematsu, S., Akira, S., Gray, D., Gray, M., MacDonald, A.S., Cunningham, A.F., Maizels,
764 R.M., 2014^{ba}. MyD88 signaling inhibits protective immunity to the gastrointestinal helminth
765 parasite *Heligmosomoides polygyrus*. J. Immunol. 193, 2984-2993.

766 ~~[Reynolds, L.A., Smith, K.A., Filbey, K.J., Harcus, Y., Hewitson, J.P., Redpath, S.A., Valdez,](#)~~
767 ~~[Y., Yebra, M.J., Finlay, B.B., Maizels, R.M., 2014b. Commensal pathogen interactions in the](#)~~
768 ~~[intestinal tract: *lactobacilli* promote infection with, and are promoted by, helminth parasites.](#)~~
769 ~~[Gut Microbes. 5, 522-532.](#)~~

770 Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S., 2011.
771 Metagenomic biomarker discovery and explanation. Genome Biol. 12.

772 Sepehri, S., Kotlowski, R., Bernstein, C.N., Krause, D.O., 2007. Microbial diversity of
773 inflamed and noninflamed gut biopsy tissues in inflammatory bowel disease. Inflamm. Bowel
774 Dis. 13, 675-683.

775 Shepherd, M.L., Swecker, W.S., Jr., Jensen, R.V., Ponder, M.A., 2012. Characterization of
776 the faecal bacteria communities of forage-fed horses by pyrosequencing of 16S rRNA V4
777 gene amplicons. FEMS Microbiol. Lett. 326, 62-68.

778 Shin, N.R., Whon, T.W., Bae, J.W., 2015. Proteobacteria: microbial signature of dysbiosis in
779 gut microbiota. Trends Biotechnol. 33, 496-503.

780 Slapeta, J., Dowd, S.E., Alanazi, A.D., Westman, M.E., Brown, G.K., 2015. Differences in
781 the faecal microbiome of non-diarrhoeic clinically healthy dogs and cats associated with
782 *Giardia duodenalis* infection: impact of hookworms and coccidia. Int. J. Parasitol. 45, 585-
783 594.

784 St-Pierre, B., Cersosimo, L.M., Ishaq, S.L., Wright, A.D., 2015. Toward the identification of
 785 methanogenic archaeal groups as targets of methane mitigation in livestock animalsr. *Front.*
 786 *Microbiol.* 6, 776.
 787 Steelman, S.M., Chowdhary, B.P., Dowd, S., Suchodolski, J., Janecka, J.E., 2012.
 788 Pyrosequencing of 16S rRNA genes in fecal samples reveals high diversity of hindgut
 789 microflora in horses and potential links to chronic laminitis. *BMC Vet. Res.* 8, 231.
 790 Stratford, C.H., McGorum, B.C., Pickles, K.J., Matthews, J.B., 2011. An update on
 791 cyathostomins: anthelmintic resistance and diagnostic tools. *Equine Vet. J. Suppl.*, 133-139.
 792 Thomsen, L.E., Petkevicius, S., Bach Knudsen, K.E., Roepstorff, A., 2005. The influence of
 793 dietary carbohydrates on experimental infection with *Trichuris suis* in pigs. *Parasitology.*
 794 131, 857-865.
 795 Traversa, D., Iorio, R., Klei, T.R., Kharchenko, V.A., Gawor, J., Otranto, D., Sparagano,
 796 O.A., 2007. New method for simultaneous species-specific identification of equine strongyles
 797 (nematoda, strongylida) by reverse line blot hybridization. *J. Clin. Microbiol.* 45, 2937-2942.
 798 Tuomisto, H., 2010. A consistent terminology for quantifying species diversity? Yes, it does
 799 exist. *Oecologia.* 164, 853-860.
 800 Uhlinger, C.A., 1991. Equine Small Strongyles - Epidemiology, Pathology, and Control.
 801 *Comp. Cont. Educ. Pract.* 13, 863-869.
 802 van Klink, R., van der Plas, F., van Noordwijk, C.G., WallisDeVries, M.F., Olf, H., 2015.
 803 Effects of large herbivores on grassland arthropod diversity. *Biol. Rev. Camb. Philos. Soc.*
 804 90, 347-366.
 805 Walk, S.T., Blum, A.M., Ewing, S.A., Weinstock, J.V., Young, V.B., 2010. Alteration of the
 806 murine gut microbiota during infection with the parasitic helminth *Heligmosomoides*
 807 *polygyrus*. *Inflamm. Bowel Dis.* 16, 1841-1849.

808 Weese, J.S., Holcombe, S.J., Embertson, R.M., Kurtz, K.A., Roessner, H.A., Jalali, M.,
 809 Wismer, S.E., 2014. Changes in the faecal microbiota of mares precede the development of
 810 post partum colic. *Equine Vet. J.* 47, 641-9.
 811 Winker, S., Woese, C.R., 1991. A definition of the domains Archaea, Bacteria and Eucarya in
 812 terms of small subunit ribosomal RNA characteristics. *Syst. Appl. Microbiol.* 14, 305-310.
 813 Woese, C.R., Gupta, R., 1981. Are archaebacteria merely derived 'prokaryotes'? *Nature.* 289,
 814 95-96.
 815 Wu, S., Li, R.W., Li, W., Beshah, E., Dawson, H.D., Urban, J.F., Jr., 2012. Worm burden-
 816 dependent disruption of the porcine colon microbiota by *Trichuris suis* infection. *PLoS One.*
 817 7, e35470.
 818
 819
 820 |

821 **Table 1: Results of faecal egg counts (FEC) analysis recorded from *Chigh* (FEC >100**
822 **eggs per gram, e.p.g.) and *Clow* (FEC <100 e.p.g.) broodmares enrolled in the study, as**
823 **well as of non-pregnant controls (NPC), over selected into two groups: 1) with mean**
824 **FEC over three consecutive samplings performed of >100 eggs per gram (e.p.g.)**
825 **(*Chigh*), and 2) with mean FEC over three consecutive samplings of <10 e.p.g. (*Clow*),**
826 **both pre-anthelmintic treatment [D0], as well as and at 2 and 14 days post-treatment**
827 **[D2, D14]. Animals Horses with FEC of > 200 e.p.g. (C200) and 0 e.p.g. (C0), are**
828 **indicated in bold.**

Group ~~i.d.~~ *Horse* Age Tapeworm Ascarid Mean FEC FEC
Animal (yrs) FEC FEC consecutive D2 D14
i.d. (e.p.g.) (e.p.g.) strongyle (e.p.g.) (e.p.g.)
(±SE) D0

<i>Chigh</i>							
	MA	6	0	0	123 (±19)	0	0
	CT	5	0	0	130 (±14)	0	0
	LE	10	0	0	113 (±17)	0	0
	PB	5	0	0	171 (±15)	0	0
	SC	4	0	0	101 (±8)	2 (±2)	0
	MSJ	6	0	0	100 (±6)	0	0
	WD	7	0	0	128 (±16)	0	0
	NS	8	0	0	120 (±11)	1 (±1)	0
	HY	7	0	0	150 (±14)	0	0
	RM	4	0	0	139 (±49)	0	0
	HS	6	0	0	200 (±39)	1 (±1)	1 (±1)
	HT	6	0	0	228 (±22)	0	0
	NO	6	0	0	271 (±17)	0	1 (±1)
	QM	4	0	0	418 (±112)	0	0
	SB	8	0	0	206 (±37)	1 (±1)	1 (±1)
	TC	4	0	0	235 (±12)	0	1 (±1)
	MQ	6	0	0	228 (±22)	0	0
	VR	4	0	0	279 (±61)	1 (±1)	0
<i>Clow</i>							
	GL	4	0	0	10 (±2)	0	0
	DWD	8	0	0	0.3 (±0.3)	0	0
	LVN	4	0	0	5 (±1)	0	0
	BB	7	0	0	1 (±1)	0	4 (±1)
	LF	5	0	0	10 (±2)	0	0
	IR	7	0	0	3 (±1)	0	0
	DDR	8	0	0	1 (±1)	0	0
	LAL	5	0	0	0.3 (±0.3)	0	0
	PT	7	0	0	0.3 (±0.3)	0	0
	PP	6	0	0	2 (±1)	0	0
	MG	5	0	0	0.5 (±0.3)	0	0
	SY	8	0	0	0	0	0
	BX	16	0	0	0	0	0
	MR	10	0	0	0	0	0

Formatted: Font: 12 pt, No underline, Font color: Auto

Formatted: Font: 12 pt, No underline, Font color: Auto

Formatted: Font: 12 pt, No underline, Font color: Auto

Formatted: Font: Italic, No underline, Font color: Auto

Formatted: Font: Italic, No underline, Font color: Auto

Formatted: Font: 12 pt, No underline, Font color: Auto

Formatted: Font: 12 pt, No underline, Font color: Auto

Formatted: Font: 12 pt, No underline, Font color: Auto

Formatted: Font: 12 pt, No underline, Font color: Auto

Formatted: Font: 12 pt, No underline, Font color: Auto

Formatted: Font: 12 pt, No underline, Font color: Auto

Formatted: Font: 12 pt, No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

Formatted: No underline, Font color: Auto

<div>Non- pregnant control</div> <div>NPC</div>	DD		0	0	0	0	0
	ED	8	0	0	0	0	0
	EP	5	0	0	0	0	0
	SWC	8	0	0	0	0	0
	BET	8	0	0	412 (±45)	-	-
	BL	12	0	0	22 (±1)	-	-
	ST	10	0	0	0	-	-

Formatted: No underline, Font color: Auto

Formatted: Font color: Text 1

Figure 1





Figure 2

A



B




Figure 3




Figure 4






Figure 5


A Chigh Clow



B C200 C0



C D0 D2 D14



D

